

Chapter 13. Wetlands

Lead Author: Scott D. Bridgham¹

Contributing Authors: J. Patrick Megonigal,² Jason K. Keller,² Carl Trettin,³ and Norman B. Bliss⁴

¹Center for Ecology and Evolutionary Biology, University of Oregon, ²Smithsonian Environmental Research Center,
³Center for Forested Wetland Research, USDA Forest Service, ⁴SAIC, USGS Center for Earth Resources
Observation and Science

KEY FINDINGS

- North America is home to approximately 41% of the global wetland area, encompassing about 2.5 million km² with a carbon pool of approximately 220 Gt, mostly in peatland soils.
 - North American wetlands currently are a CO₂ sink of approximately 70 Mt C yr⁻¹, but that estimate has an uncertainty of greater than 100%. North American wetlands are also a source of approximately 26 Mt yr⁻¹ of methane, a more potent atmospheric heat-trapping gas. The uncertainty in that flux is also greater than 100%.
 - Historically, the destruction of North American wetlands through land-use change has reduced carbon storage in wetlands by 43 Mt C yr⁻¹, primarily through the oxidation of carbon in peatland soils as they are drained and a more general reduction in carbon sequestration capacity of wetlands converted to other land uses. Methane emissions have also declined with the loss of wetland area.
 - Projections of future carbon storage and methane emissions of North American wetlands are highly uncertain and complex, but the large carbon pools in peatlands may be at risk for oxidation and release to the atmosphere as CO₂ if they become substantially warmer and drier. Methane emissions may increase with warming, but the response will likely vary with wetland type and with changes in precipitation.
 - Because of the potentially significant role of North American wetlands in methane production, the activities associated with the restoration, creation and protection of wetlands are likely to focus on the ecosystem services that wetlands provide, such as filtering of toxics, coastal erosion protection, wildlife habitat, and havens of biodiversity, rather than on carbon sequestration per se.
 - Research needs to reduce the uncertainties in carbon storage and fluxes in wetlands to provide information about management options in terms of carbon sequestration and trace gas fluxes.
-

1 INTRODUCTION

2 While there are a variety of legal and scientific definitions of a wetland (National Research Council,
3 1995; National Wetlands Working Group, 1997), most emphasize the presence of waterlogged conditions
4 in the upper soil profile during at least part of the growing season, and plant species and soil conditions
5 that reflect these hydrologic conditions. Waterlogging tends to suppress microbial decomposition more
6 than plant productivity, so wetlands are known for their ability to accumulate large amounts of carbon,
7 most spectacularly seen in large peat deposits that are often many meters deep. Thus, when examining
8 carbon dynamics, it is important to distinguish between freshwater wetlands with surface soil organic
9 matter deposits >40 cm thick (i.e., peatlands) and those with lesser amounts of soil organic matter (i.e.,
10 freshwater mineral-soil wetlands, FWMS). Some wetlands have permafrost; fluxes and pools in wetlands
11 with and without permafrost are discussed separately in Appendix 13A. We also differentiate between
12 freshwater wetlands and estuarine wetlands (salt marshes, mangroves, and mud flats) with marine-derived
13 salinity.

14 Peatlands occupy about 3% of the terrestrial global surface, yet they contain 16–33% of the total soil
15 carbon pool (Gorham, 1991; Maltby and Immirzi, 1993). Most peatlands occur between 50 and 70° N,
16 although significant areas occur at lower latitudes (Matthews and Fung, 1987; Aselmann and Crutzen,
17 1989; Maltby and Immirzi, 1993). Large areas of peatlands exist in Alaska, Canada, and in the northern
18 midwestern, northeastern, and southeastern United States (Bridgman *et al.*, 2000). This peat has formed
19 over thousands of years, and therefore the potential emissions from the large pool of soil carbon are likely
20 more significant to the global carbon budget than the current soil carbon sequestration rate. Large areas of
21 wetlands have been converted to other land uses globally and in North America (Dugan, 1993; OECD,
22 1996), which may have resulted in a net flux of carbon to the atmosphere (Armentano and Menges, 1986;
23 Maltby and Immirzi, 1993). Additionally, wetlands emit 92–237 Mt methane (CH₄) yr⁻¹, a large fraction
24 of the total annual global flux of about 600 Mt CH₄ yr⁻¹ (Ehhalt *et al.*, 2001). This is important because
25 methane is a potent greenhouse gas, second in importance to only carbon dioxide (Ehhalt *et al.*, 2001).

26 A number of previous studies have examined the role of peatlands in the global carbon balance
27 (reviewed in Mitra *et al.*, 2005). Roulet (2000) focused on the role of Canadian peatlands in the Kyoto
28 process. Here we augment these previous studies by considering all types of wetlands (not just peatlands)
29 and integrate new data to examine the carbon balance in the wetlands of Canada, the United States, and
30 Mexico.

31 Given that many undisturbed wetlands are a natural sink for carbon dioxide and a source of methane,
32 a note of caution in interpretation of our data is important. Using the International Panel on Climate
33 Change (IPCC) terminology, a radiative forcing denotes “an externally imposed perturbation in the
34 radiative energy budget of the Earth’s climate system” (Ramaswamy *et al.*, 2001). Thus, it is the change

1 from a baseline condition in greenhouse gas fluxes in wetlands that constitute a radiative forcing that will
2 impact climate change, and the emissions of greenhouse gases from unperturbed wetlands is important
3 only in establishing a baseline condition. Thus, we consider changes from historical (~1800) fluxes and
4 present and future perturbations of greenhouse gas fluxes in North American wetlands.

6 INVENTORIES

7 Current Wetland Area and Rates of Loss

8 The current and historical wetland area and rates of loss are the basis for all further estimates of pools
9 and fluxes in this chapter. The loss of wetlands has caused the oxidation of their soil carbon, particularly
10 in peatlands; reduced their ability to sequester carbon; and reduced their emissions of methane. The
11 strengths and weakness of the wetland inventories of Canada, the United States, and Mexico are discussed
12 in Appendix 13A.

13 The conterminous United States has 312,000 km² of FWMS wetlands, 93,000 km² of peatlands, and
14 23,000 km² of estuarine wetlands, which encompass 5.5% of the land area (Table 13-1). This represents
15 just 48% of the original wetland area in the conterminous United States (Table 13A-1 in Appendix 13A).
16 However, wetland losses in the United States have declined from 1,855 km² yr⁻¹ in the 1950s–1970s to
17 237 km² yr⁻¹ in the 1980s–1990s (Dahl, 2000). Such data mask large differences in loss rates among
18 wetland classes and conversion of wetlands to other classes, with potentially large effects on carbon
19 stocks and fluxes (Dahl, 2000). For example, the majority of wetland losses in the United States have
20 occurred in FWMS wetlands. As of the early 1980s, 84% of U.S. peatlands were unaltered (Armentano
21 and Menges, 1986; Maltby and Immirzi, 1993; Rubec, 1996), and, given the current regulatory
22 environment in the United States, recent rates of loss are likely small.

23
24 **Table 13-1. The area, carbon pool, net carbon balance, and methane flux from wetlands in North**
25 **America and the world.** Positive fluxes indicate net fluxes to the atmosphere, whereas negative fluxes
26 indicate net fluxes into an ecosystem. Citations and assumptions in calculations are in the text and in
27 Appendix 13A.

28
29 Canada has 1,301,000 km² of wetlands, covering 14% of its land area, of which 87% are peatlands
30 (Table 13-1). Canada has lost about 14% of its wetlands, mainly due to agricultural development of
31 FWMS wetlands (Rubec, 1996), although the ability to estimate wetland losses in Canada is limited by
32 the lack of a regular wetland inventory.

1 The wetland area in Mexico is estimated at 36,000 km² (Table 13-1), with an estimated historical loss
2 of 16,000 km² (Table 13A-1 in Appendix 13A). However, given the lack of a nationwide wetland
3 inventory and a general paucity of data, this number is highly uncertain.

4 Problems with inadequate wetland inventories are even more prevalent in lesser developed countries
5 (Finlayson *et al.*, 1999). We estimate a global wetland area of 6.0×10^6 km² (Table 13-1); thus, North
6 America currently has about 43% of the global wetland area. It has been estimated that about 50% of the
7 world's historical wetlands have been converted to other uses (Moser *et al.*, 1996).

8 9 **Carbon Pools**

10 We estimate that North American wetlands have a current soil and plant carbon pool of 220 Gt, of
11 which approximately 98% is in the soil (Table 13-1). The majority of this carbon is in peatlands, with
12 FWMS wetlands contributing about 18% of the carbon pool. The large amount of soil carbon (27 Gt) in
13 Alaskan FWMS wetlands had not been identified in previous studies (see Appendix 13A).

14 15 **Soil Carbon Fluxes**

16 North American peatlands currently have a net carbon balance of about -18 Mt C yr⁻¹ (Table 13-1),
17 but several large fluxes are incorporated into this estimate. (**Negative numbers indicate net fluxes into**
18 **the ecosystem, whereas positive numbers indicate net fluxes into the atmosphere.**) Peatlands
19 sequester -34 Mt C yr⁻¹ (Table 13A-2 in Appendix 13A), but peatlands in the conterminous United States
20 that have been drained for agriculture and forestry had a net oxidative flux of 18 Mt C yr⁻¹ as of the early
21 1980s (Armentano and Menges, 1986). Despite a substantial reduction in the rate of wetland loss since the
22 1980s (Dahl 2000), drained organic soils continue to lose carbon over many decades, so the actual flux to
23 the atmosphere is probably close to the 1980s estimate. There has also been a loss in sequestration
24 capacity in drained peatlands of 2.4 Mt C yr⁻¹ (Table 13-1), so the overall soil carbon sink of North
25 American peatlands is about 21 Mt C yr⁻¹ smaller than it would have been in the absence of disturbance.

26 Very little attention has been given to the role of FWMS wetlands in North American or global
27 carbon balance estimates, with the exception of methane emissions. Carbon sequestration associated with
28 sediment deposition is a potentially large, but poorly quantified, flux in wetlands (Stallard, 1998). Using a
29 review by Johnston (1991), we calculate a substantial carbon accumulation rate in sedimentation in
30 FWMS wetlands of -129 g C m⁻² yr⁻¹ (see Appendix 13A). However, it is extremely unlikely that the
31 actual sequestration rate is this high, as the data are probably strongly biased by researchers choosing
32 wetlands with high sediment deposition to study this process. More fundamentally, carbon in sediments
33 that are simply redistributed in the landscape due to erosion from a terrestrial source to a wetland sink
34 does not represent carbon sequestration except to the extent that decomposition rates are lower in

1 wetlands. Much of this sediment-associated carbon is probably relatively stable in upland soils, so FWMS
2 wetlands may not represent a substantial sediment carbon sink at the landscape scale. There are no data to
3 our knowledge to evaluate this important caveat. Based upon this reasoning, we somewhat arbitrarily
4 reduced our calculated FWMS wetland sediment carbon sequestration rate by 75% to -34 Mt C yr^{-1} (Table
5 13A-2 in Appendix 13A). This is still a substantial sink and an important unknown in carbon budgets. For
6 example, Stallard (1998) estimated that global wetlands are a large sediment sink, with a flux on the order
7 of -1 Gt C yr^{-1} . However, this analysis was based on many assumptions and was acknowledged by the
8 author to be a first guess at best.

9 Decomposition of soil carbon in FWMS wetlands that have been converted to other land uses appears
10 to be responsible for only a negligible loss of soil carbon currently (Table 13A-2 in Appendix 13A).
11 However, due to the historical loss of FWMS wetland area, we estimate that they currently sequester
12 21 Mt C yr^{-1} less than they did prior to disturbance (Table 13-1). This estimate has the same unknowns
13 described in the previous paragraph on current sediment carbon sequestration in FWMS wetlands.

14 We estimate that estuarine wetlands currently sequester $-9.7 \text{ Mt C yr}^{-1}$, with a historical reduction in
15 sequestration capacity of 1.4 Mt C yr^{-1} due to loss of area (Table 13-1). Despite the relatively small area
16 of estuarine wetlands, they currently contribute about 26% of total wetland carbon sequestration in the
17 conterminous United States and about 14% of the North American total. Estuarine wetlands sequester
18 carbon at a rate about 10 times higher on an area basis than other wetland ecosystems due to high
19 sedimentation rates, high soil carbon content, and constant burial due to sea level rise. Estimates of
20 sediment deposition rates in estuarine wetlands are robust, but it is unknown to what extent soil carbon
21 sequestration is divided into allochthonous carbon (sediment-derived carbon from outside the wetland)
22 and autochthonous carbon (derived from rates of plant productivity being greater than decomposition
23 within the wetland). As with FWMS wetlands, soil carbon sequestration in estuarine wetlands is
24 overestimated to the extent that allochthonous carbon simply represents redistribution of carbon in the
25 landscape. There is also large uncertainty in the area of mud flats.

26 Overall, North American wetland soils appear to be a substantial carbon sink with a net flux of
27 -70 Mt C yr^{-1} (with very large error bounds because of FWMS wetlands) (Table 13-1). The large-scale
28 conversion of wetlands to upland uses has led to a reduction in the wetland soil carbon sequestration
29 capacity of 25 Mt C yr^{-1} from the likely historical rate (Table 13-1), but this estimate is driven by large
30 losses of FWMS wetlands with their highly uncertain sedimentation carbon sink. With the current net
31 oxidative flux of 18 Mt C yr^{-1} from conterminous U.S. peatlands, we estimate that North American
32 wetlands currently sequester 43 Mt C yr^{-1} less than they did historically (Table 13A-2 in Appendix 13A).
33 Furthermore, North American peatlands and FWMS wetlands have lost 2.6 Gt and 4.9 Gt of soil carbon,
34 respectively, and collectively they have lost 2.4 Gt of plant carbon since approximately 1800. Very little

1 data exist to estimate carbon fluxes for freshwater Mexican wetlands, but because of their small area, they
2 will not likely have a large impact on the overall North American estimates.

3 The global wetland soil carbon balance has only been examined in peatlands. The current change in
4 soil carbon flux in peatlands is about 176 to 266 Mt C yr⁻¹ (Table 13A-2 in Appendix 13A), largely due to
5 the oxidation of peat drained for agriculture and forestry and secondarily due to peat combustion for fuel
6 (Armentano and Menges, 1986; Maltby and Immirzi 1993). Thus, globally peatlands are a moderate
7 atmospheric source of carbon. The cumulative historical shift in soil carbon stocks has been estimated to
8 be 5.5 to 7.1 Gt C (Maltby and Immirzi, 1993).

10 Methane and Nitrous Oxide Emissions

11 We estimate that North American wetlands emit 26 Mt CH₄ yr⁻¹ (Table 13-1). Our synthesis is
12 substantially higher than the previous estimate by Bartlett and Harriss (1993) (see Appendix 13A). A
13 mechanistic methane model yielded similar rates of 3.8 and 7.1 Mt CH₄ yr⁻¹ for Alaska and Canada,
14 respectively (Zhuang et al., 2004). For comparison, a regional inverse atmospheric modeling approach
15 estimated total methane emissions (from all sources) of 16 and 54 Mt CH₄ yr⁻¹ for boreal and temperate
16 North America, respectively (Fletcher *et al.*, 2004a).

17 Methane emissions are currently about 24 Mt CH₄ yr⁻¹ less than they were historically in North
18 American wetlands (see Table 13A-4 in Appendix 13A) because of the loss of wetland area. We do not
19 consider the effects of conversion of wetlands from one type to another (Dahl 2000), which may have a
20 significant impact on methane emissions. Similarly, we estimate that global methane emissions from
21 natural wetlands are only about half of what they were historically (Table 13A-4 in Appendix 13A).
22 However, this may be an overestimate because wetland losses have been higher in more developed
23 countries than less developed countries (Moser *et al.*, 1996), and wetlands at lower latitudes have higher
24 emissions on average (Bartlett and Harriss, 1993).

25 When we multiplied the very low published estimates of nitrous oxide emissions from natural and
26 disturbed wetlands (Joosten and Clarke, 2002) by North American wetland area, the flux was insignificant
27 (data not shown).

28 The global warming potential (GWP) of a gas depends on its instantaneous radiative forcing and its
29 lifetime in the atmosphere, with methane having GWPs of 1.9, 6.3, and 16.9 CO₂-carbon equivalents on a
30 mass basis across 500-year, 100-year, and 20-year time frames, respectively (Ramaswamy *et al.*, 2001).¹
31 Thus, depending upon the time frame and within the large confidence limits of many of our estimates in

¹ GWPs in Ramaswamy *et al.* (2001) were originally reported in CO₂-mass equivalents. We have converted them into CO₂-carbon equivalents so that the net carbon balance and methane flux columns in Table 13-1 can be directly compared by multiplying methane fluxes by the GWPs given here].

1 Table 13-1, North American wetlands as a whole currently are in a range between approximately neutral
2 and a large source of net CO₂-carbon equivalents to the atmosphere (but note caution in the *Introduction*
3 in converting this into radiative forcing). It is likely that FWMS wetlands, with their high methane
4 emissions, are a net source of CO₂-carbon equivalents to the atmosphere. In contrast, estuarine wetlands
5 are a net sink for CO₂-carbon equivalents because they support both rapid rates of carbon sequestration
6 and low methane emissions. However, caution should be exercised in using GWPs to draw conclusions
7 about changes in the net flux of CO₂-carbon equivalents because GWPs are based upon a pulse of a gas
8 into the atmosphere, whereas carbon sequestration is more or less continuous. For example, if one
9 considers continuous methane emissions and carbon sequestration in peat over time, most peatlands are a
10 net sink for CO₂-carbon equivalents because of the long lifetime of carbon dioxide sequestered as peat
11 (Frolking *et al.*, 2006).

12 13 **Plant Carbon Fluxes**

14 We estimate that wetland forests in the conterminous United States currently sequester
15 -10.3 Mt C yr⁻¹ as increased plant biomass (see Table 13A-3 in Appendix 13A). Sequestration in plants in
16 undisturbed wetland forests in Alaska and many peatlands is probably minimal, although there may be
17 substantial logging of Canadian forested peatlands that we do not have the data to account for.

18 19 **TRENDS AND DRIVERS OF WETLAND CARBON FLUXES**

20 Historically, the destruction of wetlands through land-use changes has had the largest effect on the
21 carbon fluxes and the GWPs of North American wetlands. The primary effects have been a reduction in
22 their ability to sequester carbon (a small to moderate increase in radiative forcing depending on carbon
23 sequestration by sedimentation in FWMS and estuarine wetlands), oxidation of their soil carbon reserves
24 upon drainage (a small increase in radiative forcing), and a reduction in the emission of methane to the
25 atmosphere (a moderate decrease in radiative forcing) (Table 13A-1 and Appendix 13A). While extensive
26 research has been done on carbon cycling and pools in North American wetlands, to our knowledge, this
27 is the first attempt at an overall carbon budget for all of the wetlands of North America, although others
28 have examined the carbon budget for North American peatlands as part of global assessments (Armentano
29 and Menges, 1986; Maltby and Immirzi, 1993; Joosten and Clarke, 2002). Globally, the disturbance of
30 peatlands appears to have shifted them into a net source of carbon to the atmosphere. Any positive effect
31 of wetland loss due to a reduction in their methane emissions, and hence radiative forcing, will be more
32 than negated by the loss of the many ecosystem services they provide such as havens for biodiversity,
33 recharge of groundwater, reduction in flooding, fish nurseries, etc. (Zedler and Kercher, in press).

1 A majority of the effort in examining future global change impacts on wetlands has focused on
2 northern peatlands because of their large soil carbon reserves, although under current climate conditions
3 they have modest methane emissions (Moore and Keddy, 1989; Roulet, 2000; Joosten and Clarke, 2002
4 and references therein). Data (Bartlett and Harriss, 1993; Moore *et al.*, 1998; Updegraff *et al.*, 2001) and
5 modeling (Gedney *et al.*, 2004; Zhuang *et al.*, 2004) strongly support the contention that water table
6 position and temperature are the primary environmental controls over methane emissions. How this
7 generalization plays out with future climate change is, however, more complex. For example, most
8 climate models predicted much of Canada will be warmer and drier in the future. Based upon this
9 prediction, Moore *et al.* (1998) proposed a variety of responses to climate change in the carbon fluxes
10 from different types of Canadian peatlands. Methane emissions may increase in collapsed former-
11 permafrost bogs (which will be warmer and wetter) but decrease in fens and other types of bogs (warmer
12 and drier). A methane-process model predicted that modest warming will increase global wetland
13 emissions, but larger increases in temperature will decrease emissions because of drier conditions (Cao *et al.*,
14 1998). Another methane-process model suggested that net methane emissions from northern wetlands
15 have increased by 0.08 Mt CH₄ yr⁻¹ during the twentieth century and by 1.0 Mt CH₄ yr⁻¹ during the 1980s
16 (Zhuang *et al.*, 2004). Inverse modeling also shows that atmospheric anomalies in methane during the
17 1990s may be partially explained by interannual climate effects on wetland emissions (Fletcher *et al.*,
18 2004b; Wang *et al.*, 2004). Thus, the above-mentioned studies suggest that past climate change has
19 already had an effect on wetland methane emissions and that this will only be exacerbated in the future.

20 Other important anthropogenic forcing factors that will affect future methane emissions include
21 atmospheric sulfate deposition (Vile *et al.*, 2003; Gauci *et al.*, 2004), atmospheric carbon dioxide
22 concentrations (Megonigal and Schlesinger, 1997; Vann and Megonigal, 2003), and nutrient additions
23 (Keller *et al.*, 2005). These external forcing factors in turn will interact with internal ecosystem
24 constraints such as pH and carbon quality (Moore and Roulet, 1995; Bridgham *et al.*, 1998), anaerobic
25 carbon flow (Hines and Duddleston, 2001), and net ecosystem productivity and plant community
26 composition (Whiting and Chanton, 1993; Updegraff *et al.*, 2001; Strack *et al.*, 2004) to determine the
27 actual response.

28 The effects of global change on carbon sequestration in peatlands is probably of minor importance as
29 a global flux because of the relatively low rate of peat accumulation. However, losses of soil carbon
30 stocks in peatlands drained for agriculture and forestry (Table 13A-2 in Appendix 13A) attest to the
31 possibility of large losses from the massive soil carbon deposits in northern peatlands if they become
32 substantially drier in a future climate. Furthermore, Turetsky *et al.* (2004) estimated that up to
33 5.9 Mt C yr⁻¹ are released from western Canadian peatlands by fire and predicted that increases in fire
34 frequency may cause these systems to become net atmospheric carbon sources. Northern peatlands may

1 also emit more methane with warmer temperatures, depending on changes in water table levels. The
2 effects of global change on estuarine wetlands is of concern because sequestration rates are rapid, and
3 they can be expected to increase with the rate of sea level rise provided the estuarine wetland area does
4 not decline. It remains to be determined whether rising atmospheric carbon dioxide, temperature, nitrogen
5 deposition, and shoreline construction will permit the area of estuarine wetlands to remain stable.
6

7 **OPTIONS AND MEASURES**

8 Wetland policies in the United States and Canada are driven by a variety of federal, state or
9 provincial, and local laws and regulations in recognition of the many wetland ecosystem services and
10 large historical loss rates (Lynch-Stewart *et al.*, 1999; National Research Council, 2001; Zedler and
11 Kercher, in press). Thus, any actions to enhance the ability of wetlands to sequester carbon, or reduce
12 their methane emissions, must be implemented within the context of the existing regulatory framework.
13 The most important option in the United States has already been largely achieved, and that is to reduce
14 the historical rate of peatland losses with their accompanying large oxidative losses of the stored soil
15 carbon.

16 There has been strong interest expressed in using carbon sequestration as a rationale for wetland
17 restoration and creation in the United States, Canada, and elsewhere (Wylynko, 1999; Watson *et al.*,
18 2000). However, high methane emissions from conterminous U.S. wetlands suggest that creating and
19 restoring wetlands may increase net radiative forcing, although adequate data do not exist to evaluate this.
20 Roulet (2000) came to a similar conclusion concerning the restoration of Canadian wetlands. The
21 possibility of increasing radiative forcing by creating or restoring wetlands does not apply to estuarine
22 wetlands, which emit relatively little methane compared to the carbon they sequester. Restoration of
23 drained peatlands may stop the rapid loss of their soil carbon, which may compensate for increased
24 methane emissions. However, Canadian peatlands restored from peat extraction operations increased their
25 net emissions of carbon because of straw addition during the restoration process, although it was assumed
26 that they would eventually become a net sink (Cleary *et al.*, 2005).

27 Regardless of their internal carbon balance, the area of restored wetlands is currently too small to
28 form a significant carbon sink at the continental scale. Between 1986 and 1997, only 4,157 km² of
29 uplands were converted into wetlands in the conterminous United States (Dahl, 2000). However, larger
30 areas of wetland restoration may have a significant impact on carbon sequestration. A simulation model
31 of planting 20,000 km² into bottomland hardwood trees as part of the Wetland Reserve Program in the
32 United States showed a sequestration of 4 Mt C yr⁻¹ through 2045 (Barker *et al.*, 1996), although they did
33 not account for the GWP of increased methane emissions.

1 Potentially more significant is the conversion of wetlands from one type to another; for example,
2 8.7% (37,200 km²) of the wetlands in the conterminous United States in 1997 were in a previous wetland
3 category in 1986 (Dahl, 2000). The net effect of these conversions on wetland carbon fluxes is unknown.
4 Similarly, Roulet (2000) argued that too many uncertainties exist to include Canadian wetlands in the
5 Kyoto Protocol.

6 In summary, North American wetlands form a very large carbon pool because of storage as peat and
7 are a small-to-moderate carbon sink (excluding methane effects), with the largest unknown being the role
8 of carbon sequestration by sedimentation in FWMS wetlands. With the exception of estuarine wetlands,
9 methane emissions from wetlands may largely offset any positive benefits of carbon sequestration in soils
10 and plants. Given these conclusions, it is probably unwarranted to use carbon sequestration as a rationale
11 for the protection and restoration of FWMS wetlands, although the many other ecosystem services that
12 they provide justify their protection. However, protecting and restoring peatlands will stop the loss of
13 their soil carbon (at least over the long term), and estuarine wetlands are an important carbon sink given
14 their limited areal extent and low methane emissions. The most important areas for further scientific
15 research in terms of current carbon fluxes in the United States are to establish an unbiased, landscape-
16 level sampling scheme to determine sediment carbon sequestration in FWMS and estuarine wetlands and
17 to take additional measurements of annual methane emissions to better constrain these important fluxes. It
18 would also be beneficial if the approximately decadal National Wetland Inventory (NWI) status and
19 trends data were collected in sufficient detail with respect to the Cowardin *et al.* (1979) classification
20 scheme to determine changes among mineral-soil wetlands and peatlands.

21 Canada lacks any regular inventory of its wetlands, and thus it is difficult to quantify land-use impacts
22 upon their carbon fluxes and pools. While excellent scientific data exists on most aspects of carbon
23 cycling in Canadian peatlands, Canadian FWMS and estuarine wetlands have been relatively poorly
24 studied, despite having suffered large proportional losses to land-use change. Wetland data for Mexico is
25 almost entirely lacking. Thus, anything that can be done to improve upon this would be helpful. All
26 wetland inventories should consider the area of estuarine mud flats which have the potential to sequester
27 considerable carbon.

28 Global change effects on the carbon pools and fluxes of North American wetlands are the largest
29 future unknown. We will not be able to accurately predict the role of North American wetlands as
30 potential positive or negative feedbacks to anthropogenic climate change without knowing the integrative
31 effects of changes in temperature, precipitation, atmospheric carbon dioxide concentrations, and
32 atmospheric deposition of nitrogen and sulfur within the context of internal ecosystem drivers of
33 wetlands. To our knowledge, no manipulative experiment has simultaneously measured more than two of
34 these perturbations in any North American wetland, and few have been done at any site. Modeling

1 expertise of the carbon dynamics of wetlands has rapidly improved in the last few years (Frolking *et al.*,
2 2002; Zhuang *et al.*, 2004 and references therein), but this needs even further development in the future,
3 including for FWMS wetlands.

5 ACKNOWLEDGMENTS

6 Steve Campbell [U.S. Department of Agriculture (USDA) National Resource Conservation Service
7 (NRCS), OR] synthesized the National Soil Information database so that it was useful to us. Information
8 on wetland soils within specific states was provided by Joseph Moore (USDA NRCS, AK), Robert
9 Weihrouch (USDA NRCS, WI), and Susan Platz (USDA NRCS, MN). Charles Tarnocai provided
10 invaluable data on Canadian peatlands. Thomas Dahl (U.S. Fish and Wildlife Service) explored the
11 possibility of combining NWI data with U.S. soils maps. Nigel Roulet (McGill University) gave valuable
12 advice on recent references. R. Kelman Wieder provided useful initial information on peatlands in
13 Canada.

15 REFERENCES

- 16 Armentano, T. B., and E. S. Menges, 1986: Patterns of change in the carbon balance of organic soil-wetlands of the
17 temperate zone. *Journal of Ecology*, **74**, 755–774.
- 18 Aselmann, I., and P. J. Crutzen, 1989: Global distribution of natural freshwater wetlands and rice paddies, their net
19 primary productivity, seasonality and possible methane emissions. *Journal of Atmospheric Chemistry*, **8**, 307–
20 359.
- 21 Barker, J. R., G. A. Baumgardner, D. P. Turner, and J. J. Lee, 1996: Carbon dynamics of the conservation and
22 wetland reserve program. *Journal of Soil and Water Cons.*, **51**, 340–346.
- 23 Bartlett, K. B., and R. C. Harriss, 1993: Review and assessment of methane emissions from wetlands. *Chemosphere*,
24 **26**, 261–320.
- 25 Bridgham, S. D., C.-L. Ping, J. L. Richardson, and K. Updegraff, 2000: Soils of northern peatlands: Histosols and
26 Gelisols. In: *Wetland Soils: Genesis, Hydrology, Landscapes, and Classification* (J. L. Richardson, and M. J.
27 Vepraskas, eds.), 343–370. Boca Raton, FL: CRC Press.
- 28 Bridgham, S. D., K. Updegraff, and J. Pastor, 1998: Carbon, nitrogen, and phosphorus mineralization in northern
29 wetlands. *Ecology*, **79**, 1545–1561.
- 30 Cao, M., K. Gregson, and S. Marshall, 1998: Global methane emission from wetlands and its sensitivity to climate
31 change. *Atmospheric Environment*, **32**, 3293–3299.
- 32 Cleary, J., N. T. Roulet, and T. R. Moore, 2005: Greenhouse gas emissions from Canadian peat extraction, 1990–
33 2000: A life-cycle analysis. *Ambio*, **34**, 456–461.
- 34 Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe, 1979: Classification of wetlands and deepwater habitats
35 of the United States. FWS/OBS-79/31, Washington, D.C.: Fish and Wildlife Service, U.S. Department of the
36 Interior.

- 1 Dahl, T. E., 2000: Status and Trends of Wetlands in the Conterminous United States, 1986 to 1997. Washington,
2 D.C.: Fish and Wildlife Service, U.S. Department of the Interior.
- 3 Dugan, P., ed., 1993: Wetlands in Danger—A World Conservation Atlas. New York: Oxford University Press.
- 4 Ehhalt, D., M. Prather, F. Dentener, E. Dlugokencky, E. Holland, I. Isaksen, J. Katima, V. Kirchhoff, P. Matson, P.
5 Midgley, and M. Wang, 2001: Atmospheric chemistry and greenhouse gases. In: *Climate Change 2001: The
6 Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental
7 Panel on Climate Change* (J. T. Houghton, Y. Ding, D. J. Griggs, M. Noguer, P. J. van der Linden, X. Dai, K.
8 Maskell, and C. A. Johnson, eds.), 239–287. Cambridge: Cambridge University Press.
- 9 Finlayson, C. M., N. C. Davidson, A. G. Spiers, and N. J. Stevenson, 1999: Global wetland inventory—current
10 status and future priorities. *Marine Freshwater Research*, **50**, 717–727.
- 11 Fletcher, S. E. M., P. P. Tans, L. M. Bruhwiler, J. B. Miller, and M. Heimann, 2004a: CH₄ sources estimated from
12 atmospheric observations of CH₄ and its ¹³C/¹²C isotopic ratios: 2. Inverse modeling of CH₄ fluxes from
13 geographical regions. *Global Biogeochemical Cycles*, **18**, doi:10.1029/2004GB002224.
- 14 Fletcher, S. E. M., P. P. Tans, L. M. Bruhwiler, J. B. Miller, and M. Heimann, 2004b: CH₄ sources estimated from
15 atmospheric observations of CH₄ and its ¹³C/¹²C isotopic ratios: 1. Inverse modeling of source processes. *Global
16 Biogeochemical Cycles*, **18**, doi:10.1029/2004GB002223.
- 17 Frolking, S., N. Roulet, and J. Fuglestedt, 2006: How northern peatlands influence the earth's radiative budget:
18 Sustained methane emission versus sustained carbon sequestration. *JGR-Biogeosciences*, **111**, G01008,
19 doi:01010.01029/02005JG000091.
- 20 Frolking, S., N. T. Roulet, T. R. Moore, P. M. Lafleur, J. L. Bubier, and P. M. Crill, 2002: Modeling seasonal to
21 annual carbon balance of Mer Bleue Bog, Ontario, Canada. *Global Biogeochemical Cycles*, **16**,
22 10.1029.2001GB00147, 02002.
- 23 Gauci, V., E. Matthews, N. Dise, B. Walter, D. Koch, G. Granberg, and M. Vile, 2004: Sulfur pollution suppression
24 of the wetland methane source in the 20th and 21st centuries. *Proceeding of the National Academy of Sciences,
25 USA*, **101**, 12583–12587.
- 26 Gedney, N., P. M. Cox, and C. Huntingford, 2004: Climate feedbacks from methane emissions. *Geophysical
27 Research Letters*, **31**, L20503, doi:20510.21029/22004GL020919.
- 28 Gorham, E., 1991: Northern peatlands: Role in the carbon cycle and probable responses to climatic warming.
29 *Ecological Applications*, **1**, 182–195.
- 30 Hines, M. E., and K. N. Duddleston, 2001: Carbon flow to acetate and C₁ compounds in northern wetlands.
31 *Geophysical Research Letters*, **28**, 4251–4254.
- 32 Johnston, C. A., 1991: Sediment and nutrient retention by freshwater wetlands: effects on surface water quality.
33 *Critical Reviews in Environmental Control*, **21**, 491–565.
- 34 Joosten, H., and D. Clarke, 2002: Wise Use of Mires and Peatlands - Background Principles including a Framework
35 for Decision-Making. Saarijärvi, Finland: International Mire Conservation Group and International Peat
36 Society.

- 1 Keller, J. K., S. D. Bridgham, C. T. Chapin, and C. M. Iversen, 2005: Limited effects of six years of fertilization on
2 carbon mineralization dynamics in a Minnesota fen. *Soil Biology and Biochemistry*, **37**, 1197–1204.
- 3 Lynch-Stewart, P., I. Kessel-Taylor, and C. Rubec, 1999: Wetlands and Government: Policy and Legislation for
4 Wetland Conservation in Canada, No. 1999-1: North American Wetlands Conservation Council (Canada).
- 5 Maltby, E., and P. Immirzi, 1993: Carbon dynamics in peatlands and other wetland soils, regional and global
6 perspectives. *Chemosphere*, **27**, 999–1023.
- 7 Matthews, E., and I. Fung, 1987: Methane emission from natural wetlands: Global distribution, area, and
8 environmental characteristics of sources. *Global Biogeochemical Cycles*, **1**, 61–86.
- 9 Megonigal, J. P., and W. H. Schlesinger, 1997: Enhanced CH₄ emissions from a wetland soil exposed to elevated
10 CO₂. *Biogeochemistry*, **37**, 77–88.
- 11 Mitra, S., R. Wassmann, and P. L. G. Vlek, 2005: An appraisal of global wetland area and its organic carbon stock.
12 *Current Science*, **88**, 25–35.
- 13 Moore, D. R. J., and P. A. Keddy, 1989: The relationship between species richness and standing crop in wetlands:
14 the importance of scale. *Vegetatio*, **79**, 99–106.
- 15 Moore, T. R., and N. T. Roulet, 1995: Methane emissions from Canadian peatlands. In: *Soils and Global Change*
16 (R. Lal, J. Kimble, E. Levine, and B. A. Stewart, eds.), 153–164. Boca Raton, FL: Lewis Publishers.
- 17 Moore, T. R., N. T. Roulet, and J. M. Waddington, 1998: Uncertainty in predicting the effect of climatic change on
18 the carbon cycling of Canadian peatlands. *Climatic Change*, **40**, 229–245.
- 19 Moser, M., C. Prentice, and S. Frazier, 1996, A global overview of wetland loss and degradation: Ramsar 6th
20 Meeting of the Conference of the Contracting Parties in Brisbane, Australia.
- 21 National Research Council, 1995: Wetlands: Characteristics and Boundaries. Washington, D.C.: National Academy
22 Press.
- 23 National Research Council, 2001: Compensating for wetland losses under the clean water act. Washington, D.C.:
24 National Academy Press.
- 25 National Wetlands Working Group, 1997: The Canadian Wetland Classification System. Waterloo, Ontario,
26 Canada: Wetlands Research Centre, University of Waterloo.
- 27 OECD, 1996: Guidelines for aid agencies for improved conservation and sustainable use of tropical and subtropical
28 wetlands. Paris: Organization for Economic Co-operation and Development.
- 29 Ramaswamy, V., O. Boucher, J. Haigh, D. Hauglustaine, J. Haywood, G. Myhre, T. Nakajima, G. Y. Shi, and
30 S. Solomon, 2001: Radiative forcing of climate change. In: *Climate Change 2001: The Scientific Basis.*
31 *Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate*
32 *Change* (J. T. Houghton, Y. Ding, D. J. Griggs, M. Noguer, P. J. van der Linden, X. Dai, K. Maskell, and C. A.
33 Johnson, eds.), 349–416. Cambridge: Cambridge University Press.
- 34 Roulet, N. T., 2000: Peatlands, carbon storage, greenhouse gases, and the Kyoto Protocol: prospects and
35 significance for Canada. *Wetlands*, **20**, 605–615.
- 36 Rubec, C., 1996: The status of peatland resources in Canada. In: *Global Peat Resources*. E. Lappalainen, ed., 243–
37 252. Jyskä, Finland: International Peat Society and Geological Survey of Finland.

- 1 Stallard, R. F., 1998: Terrestrial sedimentation and the carbon cycle: Coupling weathering and erosion to carbon
2 burial. *Global Biogeochemical Cycles*, **12**, 231–257.
- 3 Strack, M., J. M. Waddington, and E.-S. Tuittila, 2004: Effect of water table drawdown on northern peatland
4 methane dynamics: Implications for climate change. *Global Biogeochemical Cycles*, **18**, GB4003,
5 doi:4010.1029/2003GB002209, 002004.
- 6 Turetsky, M. R., B. D. Amiro, E. Bosch, and J. S. Bhatti, 2004: Historical burn area in western Canadian peatlands
7 and its relationship to fire weather indices. *Global Biogeochemical Cycles*, **18**, GB4014,
8 doi:1029/2004GB002222, 002004.
- 9 Updegraff, K., S. D. Bridgman, J. Pastor, P. Weishampel, and C. Harth, 2001: Response of CO₂ and CH₄ emissions
10 in peatlands to warming and water-table manipulation. *Ecological Applications*, **11**, 311–326.
- 11 Vann, C. D., and J. P. Megonigal, 2003: Elevated CO₂ and water depth regulation of methane emissions:
12 Comparison of woody and non-woody wetland plant species. *Biogeochemistry*, **63**, 117–134.
- 13 Vile, M. A., S. D. Bridgman, R. K. Wieder, and M. Novák, 2003: Atmospheric sulfur deposition alters pathways of
14 gaseous carbon production in peatlands. *Global Biogeochemical Cycles*, **17**, 1058–1064.
- 15 Wang, J. S., J. A. Logan, M. B. McElroy, B. N. Duncan, I. A. Megretskaia, and R. M. Yantosca, 2004: A 3-D model
16 analysis of the slowdown and interannual variability in the methane growth rate from 1988 to 1997. *Global
17 Biogeochemical Cycles*, **18**, GB3011, doi:101029/102003GB002180.
- 18 Watson, R. T., I. R. Noble, B. Bolin, N. H. Ravindranath, D. J. Verardo, and D. J. Dokken, 2000: IPCC Special
19 Report on Land Use, Land-Use Change and Forestry. Cambridge, UK: Cambridge University Press.
- 20 Whiting, G. J., and J. P. Chanton, 1993: Primary production control of methane emissions from wetlands. *Nature*,
21 **364**, 794–795.
- 22 Wylyenko, D., ed., 1999: Prairie wetlands and carbon sequestration: assessing sinks under the Kyoto Protocol.
23 Winnipeg, Manitoba, Canada: Institute for Sustainable Development, Ducks Unlimited Canada, and Wetlands
24 International.
- 25 Zedler, J. B., and S. Kercher, in press: Wetland resources: status, trends, ecosystem services, and restorability.
26 *Annual Review of Environmental Resources*.
- 27 Zhuang, Q., J. M. Melillo, D. W. Kicklighter, R. G. Prin, A. D. McGuire, P. A. Steudler, B. S. Felzer, and S. Hu,
28 2004: Methane fluxes between terrestrial ecosystems and the atmosphere at northern high latitudes during the
29 past century: A retrospective analysis with a process-based biogeochemistry model. *Global Biogeochemical
30 Cycles*, **18**, GB 3010, doi:3010.1029/2004GB002239.

1 **Table 13-1. The area, carbon pool, net carbon balance, and methane flux from wetlands in North America and the world.** Positive fluxes indicate net
 2 fluxes to the atmosphere, whereas negative fluxes indicate net fluxes into an ecosystem. Citations and assumptions in calculations are in the text and in Appendix
 3 13A.

	Area ^a (km ²)		Carbon Pool ^b (Gt C)		Net Carbon Balance ^c (Mt C yr ⁻¹)		Historical Loss in Sequestration Capacity (Mt C yr ⁻¹)		Methane Flux (Mt CH ₄ yr ⁻¹)	
Canada										
Peatland	1,135,608	****	149	****	-19	***	0.3	*	3.2	**
Freshwater Mineral	158,720	**	4.9	**	-5.1	*	6.5	*	5.7	*
Estuarine	6,400	***	0.1	***	-1.3	**	0.5	*	0.0	***
Total	1,300,728	****	154	****	-25	**	7.2	*	8.9	*
Alaska										
Peatland	132,196	****	15.9	**	-2.0	**	0.0	****	0.3	*
Freshwater Mineral	555,629	****	27.1	**	-18	*	0.0	****	1.4	*
Estuarine	8,400	****	0.1	***	-1.9	**	0.0	****	0.1	***
Total	696,224	****	43.2	**	-22	*	0.0	****	1.8	*
Conterminous United States										
Peatland	93,477	****	14.4	***	4	*	2.1	*	3.4	**
Freshwater Mineral	312,193	****	6.2	***	-18	*	15	*	11.2	**
Estuarine	23,000	****	0.6	****	-4.9	**	0.4	*	0.1	***
Total	428,670	****	21.2	***	-19	*	17	*	14.7	**
U.S. Total	1,124,895	****	64	**	-41	*	17	*	17	**
Mexico										
Peatland	10,000	*	1.5	*	-1.6	*	ND ^d	*	0.4	*
Freshwater Mineral	20,685	*	0.4	*	-0.7	*	ND	*	0.7	*
Estuarine	5,000	*	0.2	*	-1.6	*	0.5	*	0.0	*
Total	35,685	*	2.1	*	-3.9	*	ND	*	1.1	*
North America										
Peatland	1,371,281	****	180	****	-18	*	2.4	*	7	**

Freshwater Mineral	1,047,227	****	39	***	-42	*	21	*	19	*
Estuarine	42,800	***	1.0	***	-9.7	**	1.4	*	0.2	**
Total	2,461,308		220		-70	*	25	*	26	*
Global										
Peatland	3,443,000	***	460	***	150	**	16	*	37	**
Freshwater Mineral	2,315,000	***	46	***	-75	*	87	*	68	**
Estuarine	203,000	*	5.4	*	-43	*	13.2	*	1.5	**
Total	5,961,000	***	511	***	32	*	116	*	107	**

1
2
3
4
5
6
7
8
9
10
11
12
13
14

^aEstuarine includes salt marsh, mangrove, and mudflat, except for Mexico and global for which no mudflat estimates were available.

^bIncludes soil C and plant C, but overall soil C is 98% of the total pool.

^cIncludes soil C sequestration, plant C sequestration, and loss of C due to drainage of wetlands. Plant C sequestration and soil oxidative flux due to drainage are either unknown or negligible for North American wetlands except for the conterminous United States (see Appendix 13A).

^dNo data.

The error categories are as follows:

***** = 95% certain that the actual value is within 10% of the estimate reported.

**** = 95% certain that the actual value is within 25%.

*** = 95% certain that the actual value is within 50%.

** = 95% certain that the actual value is within 100%.

* = uncertainty > 100%

Appendix 13A

Wetlands – Supplemental Material

INVENTORIES

Current Wetland Area and Rates of Loss

The ability to estimate soil carbon pools and fluxes in North American wetlands is constrained by the national inventories (or lack thereof) for Canada, the United States, and Mexico (Davidson *et al.*, 1999). The National Wetland Inventory (NWI) program of the United States has repeatedly sampled several thousand wetland sites using aerial photographs and more limited field verification. The data are summarized in a series of reports detailing changes in wetland area in the conterminous United States for the periods of the mid-1950s to mid-1970s (Frayer *et al.*, 1983), mid-1970s to mid-1980s (Dahl and Johnson, 1991), and 1986 to 1997 (Dahl, 2000). We used these relatively high-quality data sets extensively for estimating wetland area and loss rates in the conterminous United States, including mud flats. However, the usefulness of the NWI inventory reports for carbon budgeting was limited by the level of classification used to define wetland categories with the Cowardin *et al.* (1979) wetland classification system. At the level used in the national status and trend reports, vegetated freshwater wetlands are classified by dominant physiognomic vegetation type, and it is impossible to make the important distinction between wetlands with deep organic soils (i.e., peatlands) and wetlands with mineral soils. The data are not at an adequate spatial resolution to combine with U.S. Department of Agriculture (USDA) National Resources Conservation Service (NRCS) soil maps to discriminate between the two types of wetlands (T. Dahl, personal comm.). Because of these data limitations, we used the NRCS soil inventory of peatlands (i.e., Histosols and Histels, or peatlands with and without permafrost, respectively) to estimate historical peatland area (Bridgham *et al.*, 2000) and combined these data with regional estimates of loss (Armentano and Menges, 1986) to estimate current peatland area in the conterminous United States. We calculated the current area of freshwater mineral-soil (FWMS) wetlands in the conterminous United States by subtracting peatland area from total wetland area (Dahl, 2000). This approach was limited by the Armentano and Menges peatland area data being current only up to the early 1980s, although large losses of peatlands since then are unlikely due to the institution of wetland protection laws.

We used a similar approach for Alaskan peatlands: peatland area was determined by the NRCS soil inventory [N. Bliss, query of the NRCS State Soil Geographic (STATSGO) database, February 2006] and overall wetland inventory was determined by standard NWI methods (Hall *et al.*, 1994). However, our

1 peatland estimate of 132,000 km² (Table 13A-1) is 22% of the often cited value by Kivinen and Pakarinen
2 (1981) of 596,000 km².

3
4 **Table 13A-1. Current and historical area of wetlands in North America and the world (×10³ km²).**
5 Historical refers to approximately 1800, unless otherwise specified
6

7 Kivinen and Pakarinen also used NRCS soils data (Rieger *et al.*, 1979) for their peatland estimates, but
8 they defined a peatland as having a minimum organic layer thickness of 30 cm, whereas the current U.S.
9 and Canadian soil taxonomies require a 40-cm thickness. The original 1979 Alaska soil inventory has
10 been reclassified with current U.S. soil taxonomy (J. Moore, Alaska State Soil Scientist, personal comm.).
11 Using the reclassified soil inventory, Alaska has 417,000 km² of wetlands with a histic modifier that are
12 not Histosols or Histels, indicating significant carbon accumulation in the surface horizons of FWMS
13 wetlands. Thus, we conclude that Kivinen and Pakarinen's Alaska peatland area estimate is higher
14 because many Alaskan wetlands have a thin organic horizon that is not deep enough to qualify as a
15 peatland under current soil taxonomy. Our smaller peatland area significantly lowers our estimate of
16 carbon pools and fluxes in Alaskan peatlands compared to earlier studies (see *Carbon Pools* below).

17 A regular national inventory of Canada's wetlands has not been undertaken, although wetland area
18 has been mapped by ecoregion (National Wetlands Working Group, 1988). Extensive recent effort has
19 gone into mapping Canadian peatlands (Tarnocai, 1998; Tarnocai *et al.*, 2005). We calculated mineral-
20 soil wetlands as the difference between total wetland area and peatland area in National Wetland Working
21 Group (1988). Historical FWMS wetland area was obtained from Rubec (1996). There are no reliable
22 country-wide estimates of mud flat area for Canada, but a highly uncertain extrapolation from a limited
23 number of regional estimates was possible.

24 No national wetland inventories have been done for Mexico. Current freshwater wetland estimates for
25 Mexico were taken from Davidson *et al.* (1999), who used inventories of discrete wetland regions
26 performed by a variety of organizations. Thus, freshwater wetland area estimates for Mexico are highly
27 unreliable and are possibly a large underestimate. For salt marshes and mangroves area in Mexico, we
28 used the estimates compiled by Mendelsohn and McKee (2000), which are similar to estimates reported
29 in Davidson *et al.* (1999) and Spalding *et al.* (1997). There are no reliable estimates of mud flat area for
30 Mexico.

31 **CARBON POOLS**

32 **Freshwater Mineral-Soil (Gleysol) Carbon Pools**

33 Gleysol is a soil classification used by the Food and Agriculture Organization (FAO) and many
34 countries that denotes mineral soils formed under waterlogged conditions (FAO-UNESCO, 1974).
35

1 Tarnocai (1998) reported a soil carbon density of 200 Mg C ha⁻¹ for Canadian Gleysols but did not
2 indicate to what depth this extended. Batjes (1996) determined soil carbon content globally from the *Soil*
3 *Map of the World* (FAO, 1991) and a large database of soil pedons. He gave a very similar average value
4 for soil carbon density of 199 Mg C ha⁻¹ (CV² = 212%, n = 14 pedons) for Gleysols of the world to 2-m
5 depth; to 1-m depth, he reported a soil carbon density of 131 Mg C ha⁻¹ (CV = 109%, n = 142 pedons).

6 Gleysols are not part of the U.S. soil taxonomy scheme, and mineral soils with attributes reflecting
7 waterlogged conditions are distributed among numerous soil groups. We used the NRCS State Soil
8 Geographic (STATSGO) soils database to query for soil carbon density in “wet” mineral soils of the
9 conterminous United States (all soils that had a surface texture described as peat, muck, or mucky peat, or
10 appeared on the 1993 list of hydric soils, which were not classified as Histosols) (N. Bliss, query of
11 NRCS STATSGO database, Dec. 2005). We found soil carbon densities of 162 Mg C ha⁻¹ for FWMS
12 wetlands in the conterminous United States and Mexico, which was used in this analysis.

13 However, some caution is necessary regarding the use of Gleysol or wet mineral soil carbon densities,
14 as apparently they include large areas of seasonally wet soils that are not considered wetlands by the more
15 conservative definition of wetlands used by the United States and many other countries and organizations.
16 For example, Eswaran *et al.* (1995) estimated that global wet mineral-soil area was 8,808,000 km², which
17 is substantially higher than the commonly accepted mineral-soil wetland area estimated by Matthews and
18 Fung (1987) of 2,289,000 km² and Aselmann and Crutzen (1989) of 2,341,000 km², even accounting for
19 substantial global wetland loss. In our query of the NRCS STATSGO database for the United States, we
20 found 1,258,000 km² of wet soils in the conterminous United States versus our estimate of 312,000 km²
21 of FWMS wetlands currently and 762,000 km² historically (Table 13A-1). We assume that including
22 these wet-but-not-wetland soils will decrease the estimated soil carbon density, but to what degree we do
23 not know. However, just considering the differences in area will give large differences in the soil carbon
24 pool. For example, Eswaran *et al.* (1995) estimated that wet mineral soils globally contain 108 Gt C to
25 1-m depth, whereas our estimate is 46 Gt C to 2-m depth (Table 13A-2).

26 For Alaska, many soil investigations have been conducted since the STATSGO soil data was coded.
27 We updated STATSGO by calculating soil carbon densities from data obtained from the NRCS on
28 479 pedons collected in Alaska, and then we used this data for both FWMS wetlands and peatlands. For
29 some of the Histosols, missing bulk densities were calculated using averages of measured bulk densities
30 for the closest matching class in the USDA Soil Taxonomy (NRCS, 1999). A matching procedure was
31 developed for relating sets of pedons to sets of STATSGO components. If there were multiple
32 components for each map unit in STATSGO, the percentage of the component was used to scale area and
33 carbon data. We compared matching sets of pedons to sets of components at the four top levels of the

² CV is the “coefficient of variation,” or 100 times the standard deviation divided by the mean.

1 U.S. Soil Taxonomy: Orders, Suborders, Great Groups, and Subgroups. For example, the soil carbon for
2 all pedons having the same soil order were averaged, and the carbon content was applied to all of the soil
3 components of the same order (e.g., Histosol pedons are used to characterize Histosol components). At
4 the Order level, all components were matched with pedon data. At the suborder level, pedon data were not
5 available to match approximately 20,000 km² (compared to the nearly 1,500,000-km² area of soil in the
6 state), but the soil characteristics were more closely associated with the appropriate land areas than at the
7 Order level. At the Great Group and Subgroup levels, pedon data were unavailable for much larger areas,
8 even though the quality of the data when available became better. For this study, we used the Suborder-
9 level matching. The resulting soil carbon density for Alaskan FWMS wetlands was 469 Mg C ha⁻¹,
10 reflecting large areas of wetlands with a histic epipedon as noted above.

11 12 **Peatland Soil Carbon Pools**

13 The carbon pool of permafrost and non-permafrost peatlands in Canada had been previously
14 estimated by Tarnocai *et al.* (2005) based upon an extensive database. Good soil-carbon density data are
15 unavailable for peatlands in the United States, as the NRCS soil pedon information typically only goes to
16 a maximum depth of between 1.5 to 2 m, and many peatlands are deeper than this. Therefore, we used the
17 carbon density estimates of Tarnocai *et al.* (2005) of 1,441 Mg C ha⁻¹ for Histosols and 1,048 Mg C ha⁻¹
18 for Histels to estimate the soil carbon pool in Alaskan peatlands.

19 The importance of our using a smaller area of Alaskan peatlands becomes obvious here. Using the
20 larger area from Kivinen and Pakarinen (1981), Halsey *et al.* (2000) estimated that Alaskan peatlands
21 have a soil carbon pool of 71.5 Gt, almost 5-fold higher than our estimate. However, some of the
22 difference in soil carbon between the two estimates can be accounted for by the 26 Gt C that we
23 calculated resides in Alaskan FWMS wetlands (Table 13A-2).

24 25 **Table 13A-2. Soil carbon pools (Gt) and fluxes (Mt yr⁻¹) of wetlands in North America and the world.**

26 “Sequestration in current wetlands” refers to carbon sequestration in wetlands that currently exist;
27 “oxidation in former wetlands” refers to emissions from wetlands that have been converted to non-wetland
28 uses or conversion among wetland types due to human influence; “historical loss in sequestration capacity”
29 refers to the loss in the carbon sequestration function of wetlands that have been converted to non-wetland
30 uses; “change in flux from wetland conversions” is the sum of the two previous fluxes. Positive flux
31 numbers indicate a net flux into the atmosphere, whereas negative numbers indicate a net flux into the
32 ecosystem

33
34 The peatlands of the conterminous United States are different in texture, and probably depth, from those
35 in Canada and Alaska, so it is probably inappropriate to use the soil carbon densities for Canadian

1 peatlands for those in the conterminous United States. For example, we compared the relative percentage
2 of the Histosol suborders (excluding the small area of Folists, as they are predominantly upland soils) for
3 Canada (Tarnocai, 1998), Alaska (updated STATSGO data, J. Moore, personal comm.), and the
4 conterminous U.S. (NRCS, 1999). The relative percentage of Fibrists, Hemists, and Saprists, respectively,
5 in Canada are 37%, 62%, and 1%, in Alaska are 53%, 27%, and 20%, and in the conterminous United
6 States are 1%, 19%, and 80%. Using the STATSGO database (N. Bliss, query of NRCS STATSGO
7 database, December 2005), the average soil carbon density for Histosols in the conterminous United
8 States is 1,089 Mg C ha⁻¹, but this is an underestimate as many peatlands were not sampled to their
9 maximum depth. Armentano and Menges (1986) reported average carbon density of conterminous U.S.
10 peatlands to 1-m depth of 1,147 to 1,125 Mg C ha⁻¹. Malterer (1996) gave soil carbon densities of
11 conterminous U.S. peatlands of 2,902 Mg C ha⁻¹ for Fibrist, 1,874 Mg C ha⁻¹ for Hemists, and 2,740 Mg
12 C ha⁻¹ for Saprists, but it is unclear how he derived these estimates. Batjes (1996) and Eswaran *et al.*
13 (1995) gave average soil carbon densities to 1-m depth for global peatlands of 776 and 2,235 Mg C ha⁻¹,
14 respectively. We chose to use an average carbon density of 1,500 Mg C ha⁻¹, which is in the middle of the
15 reported range.

16

17 **Estuarine Soil Carbon Pools**

18 Tidal wetland soil carbon density was based on a country-specific analysis of data reported in an
19 extensive compilation by Chimura *et al.* (2003). There were more observations for the United States
20 (n = 75) than Canada (n = 34) or Mexico (n = 4), and consequently there were more observations of
21 marshes than mangroves. The Canadian salt marsh estimate was used for Alaska, and country-specific
22 marsh or mangrove estimates were used for mudflats. Although Chimura *et al.* (2003) reported some
23 significant correlations between soil carbon density and mean annual temperature, scatter plots suggest
24 the relationships are weak or driven by a few sites. Thus, we did not separate the data by region or latitude
25 and used mean values for scaling. Chimura *et al.* (2003) assumed a 50-cm-deep profile for the soil carbon
26 pool, which may be an underestimate.

27

28 **Plant Carbon Pools**

29 While extensive data on plant biomass in individual wetlands have been published, no systematic
30 inventory of wetland plant biomass has been undertaken in North America. Nationally, the forest carbon
31 biomass pool (including aboveground and belowground biomass) has been estimated to be 5.49 kg C m⁻²
32 (Birdsey, 1992), which we used for forested wetlands in the United States and Canada. This approach
33 assumes that wetland forests do not have substantially different biomass carbon densities from upland
34 forests. There is one regional assessment of forested wetlands in the southeastern United States, which

1 comprise approximately 35% of the total forested wetland area in the conterminous United States. We
2 utilized the southeastern U.S. regional inventory to evaluate this assumption; aboveground tree biomass
3 averaged $125.2 \text{ m}^3 \text{ ha}^{-1}$ for softwood stands and $116.1 \text{ m}^3 \text{ ha}^{-1}$ for hardwood stands. Using an average
4 wood density and carbon content, the carbon density for these forests would be 3.3 kg C m^{-2} for softwood
5 stands and 4.2 kg C m^{-2} for hardwood stands. However, these estimates do not include understory
6 vegetation, belowground biomass, or dead trees, which account for 49% of the total forest biomass
7 (Birdsey, 1992). Using that factor to make an adjustment for total forest biomass, the range would be 4.9
8 to 6.6 kg C m^{-2} for the softwood and hardwood stands, respectively. Accordingly, the assumption of using
9 5.49 kg C m^{-2} seems reasonable for a national-level estimate.

10 The area of forested wetlands in Canada came from Tarnocai *et al.* (2005), for Alaska from Hall *et al.*
11 (1994), and for the conterminous United States from Dahl (2000).

12 Since Tarnocai *et al.* (2005) divided Canadian peatland area into bog and fen, we used aboveground
13 biomass for each community type from Vitt *et al.* (2000), and assumed that 50% of biomass is
14 belowground. We used the average bog and fen plant biomass from Vitt *et al.* (2000) for Alaskan
15 peatlands. For other wetland areas, we used an average value of $2,000 \text{ g C m}^{-2}$ for non-forested wetland
16 biomass carbon density (Gorham, 1991).

17 Tidal marsh root and shoot biomass data were estimated from a compilation in Table 8-7 in Mitsch
18 and Gosselink, (1993). There was no clear latitudinal or regional pattern in biomass, so we used mean
19 values for each. Mangrove biomass has been shown to vary with latitude (Twilley *et al.*, 1992). Biomass
20 was estimated from an empirical equation for aboveground biomass as a function of latitude (Twilley *et*
21 *al.* 1992). We made a simple estimate using a single latitude that visually bisected the distribution of
22 mangroves either in the United States (26.9°) or Mexico (23.5°). Total biomass was estimated using a
23 root-to-shoot ratio of 0.82 and a carbon-mass-to-biomass ratio of 0.45, both from Twilley *et al.* (1992).

24 Plant biomass carbon data are presented in Table 13A-3.

25
26 **Table 13A-3. Plant carbon pools (Gt) and fluxes (Mt yr^{-1}) of wetlands in North America and the**
27 **world.** Positive flux numbers indicate a net flux into the atmosphere, whereas negative numbers indicate a
28 net flux into the ecosystem
29

30 CARBON FLUXES

31 Peatland Soil Carbon Accumulation Rates

32 Most studies report the long-term apparent rate of carbon accumulation (LORCA) in peatlands based
33 upon basal peat dates, but this assumes a linear accumulation rate through time. However, due to the slow
34 decay of the accumulated peat, the true rate of carbon accumulation will always be less than the LORCA

1 (Clymo *et al.*, 1998), so most reported rates are inherently biased upwards. Tolonen and Turunen (1996)
2 found that the true rate of peat accumulation was about 67% of the LORCA.

3 For estimates of soil carbon sequestration in conterminous U.S. peatlands, we used the data from 82
4 sites and 215 cores throughout eastern North America (Webb and Webb III, 1988). They reported a
5 median accumulation rate of 0.066 cm yr⁻¹ (mean = 0.092, sd = 0.085). We converted this value into a
6 carbon accumulation rate of -1.2 Mg C ha⁻¹ yr⁻¹ by assuming 58% C (see NRCS Soil Survey Laboratory
7 Information Manual, available on-line at <http://soils.usda.gov/survey/nscd/lim/>), a bulk density of 0.59 g
8 cm⁻³, and an organic matter content of 55%. (Positive carbon fluxes indicate net fluxes to the atmosphere,
9 whereas negative carbon fluxes indicate net fluxes into an ecosystem.) The bulk density and organic
10 matter content were the average from all Histosol soil map units greater than 202.5 ha (n = 5,483) in the
11 conterminous United States from the National Soil Information System (NASIS) data base provided by S.
12 Campbell (USDA NRCS, Portland, OR). For comparison, Armentano and Menges (1986) used soil
13 carbon accumulation rates that ranged from -0.48 Mg C ha⁻¹ yr⁻¹ in northern conterminous U.S. peatlands
14 to -2.25 Mg C ha⁻¹ yr⁻¹ in Florida peatlands.

15 Peatlands accumulate lesser amounts of soil carbon at higher latitudes, with especially lower rates
16 occurring in permafrost peatlands (Ovenden, 1990, Robinson and Moore, 1999). The rates used in this
17 report reflect this gradient, going from -0.13 to -0.19 to -1.2 Mg C ha⁻¹ yr⁻¹ in permafrost peatlands, non-
18 permafrost Canadian and Alaskan peatlands, and peatlands in the conterminous United States and
19 Mexico, respectively (Table 13A-2).

21 **Freshwater Mineral-Soil Wetland Carbon Accumulation Rates**

22 Many studies have estimated sediment deposition rates in FWMS wetlands, with an average rate of
23 1,680 g m⁻² yr⁻¹ (range 0 to 7,840) in a review by Johnston (1991). Assuming 7.7% carbon for FWMS
24 wetlands (Batjes, 1996), this gives a substantial accumulation rate of -129 g C m⁻² yr⁻¹. Johnston (1991)
25 found many more studies that just reported vertical sediment accumulation rates, with an average of
26 0.69 cm yr⁻¹ (range -0.6 to 2.6). If we assume a bulk density of 1.38 g cm⁻³ for FWMS wetlands (Batjes,
27 1996), this converts into an impressive accumulation rate of -733 g C m⁻² yr⁻¹. However, we believe that
28 these values cannot be used directly as estimates of carbon sequestration rates for of two reasons. First, it
29 is likely that researchers preferentially choose wetlands with high sedimentation rates to study this
30 process. Secondly, and more fundamentally, at a landscape scale a redistribution of sediments from
31 uplands to wetlands represents no net carbon sequestration if the decomposition rate of carbon is the same
32 in both environments. The carbon associated with sediments is likely relatively recalcitrant and often
33 physically protected from decomposers by association with mineral soils. Thus, despite the anaerobic
34 conditions in wetlands, decomposition rates in deposited sediments may not be substantially lower than in

1 the uplands from which those sediments were eroded. Because of this reasoning, we somewhat arbitrarily
2 reduced our calculated rates of carbon sequestration in FWMS wetlands by 75% to $-34 \text{ g C m}^{-2} \text{ yr}^{-1}$, which
3 still represents a substantial carbon sink.

4 Agriculture typically increases sedimentation rates by 10- to 100-fold, and 90% of sediments are
5 stored within the watershed, or about 3 Gt yr^{-1} in the United States (Meade *et al.*, 1990, as cited in
6 Stallard, 1998). Converting this to 1.5% C equates to -45 Mt C yr^{-1} , part of which will be stored in
7 wetlands and is well within our estimated storage rate in FWMS wetlands (Table 13A-2).

9 **Estuarine Carbon Accumulation Rates**

10 Carbon accumulation in tidal wetlands was assumed to be entirely in the soil pool. This should
11 provide a reasonable estimate because marshes are primarily herbaceous, and mangrove biomass should
12 be in steady state unless the site was converted to another use. An important difference between soil
13 carbon sequestration in tidal and non-tidal systems is that tidal sequestration occurs primarily through
14 burial driven by sea level rise. For this reason, carbon accumulation rates can be estimated well with data
15 on changes in soil surface elevation and carbon density. Rates of soil carbon accumulation were
16 calculated from Chimura *et al.* (2003) as described for the soil carbon pool (above). These estimates are
17 based on a variety of methods, such as ^{210}Pb dating and soil elevation tables, which integrate vertical soil
18 accumulation rates over periods of time ranging from 1–100 yr.

20 **Extractive Uses of Peat**

21 Use of peat for energy production is, and always has been, negligible in North America, as opposed to
22 other parts of the world (WEC, 2001). However, Canada produces a greater volume of horticultural and
23 agricultural peat than any other country in the world (WEC, 2001). Currently, 124 km^2 of Canadian
24 peatlands have been under extraction now or in the past (Cleary *et al.*, 2005). A life-cycle analysis by
25 these authors estimated that as of 1990 Canada emitted 0.9 Mt yr^{-1} of $\text{CO}_2\text{-C}$ equivalents through peat
26 extraction. The U.S. production of horticultural peat is about 19% of Canada's (Joosten and Clarke,
27 2002), which assuming a similar life-cycle as for Canada, suggests that the United States produces 0.2 Mt
28 of $\text{CO}_2\text{-C}$ equivalents through peat extraction.

30 **Methane Fluxes**

31 Moore *et al.* (1995) reported a range of methane fluxes from 0 to $130 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$ from 120
32 peatland sites in Canada, with the majority $<10 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$. They estimated a low average flux rate of
33 2 to $3 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$, which equaled an emission of 2–3 $\text{Mt CH}_4 \text{ yr}^{-1}$ from Canadian peatlands. We used
34 an estimate of $2.5 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$ for Canadian peatlands and Alaskan freshwater wetlands (Table 13A-4).

1
2 **Table 13A-4. Methane fluxes (Mt yr^{-1}) from wetlands in North America and the world.**

3
4 To our knowledge, the last synthesis of methane fluxes was done by Bartlett and Harriss (1993). We
5 supplemented their analysis with all other published field studies (using chamber or eddy covariance
6 techniques) we could find that reported annual or average daily methane fluxes in the conterminous
7 United States (Table 13A-5). We excluded a few studies that used cores or estimated diffusive fluxes.

8
9 **Table 13A-5. Methane fluxes measured in the conterminous United States.** The conversion factor is the
10 ratio of the daily average flux to the measured annual flux $\times 10^3$. The calculated annual flux was
11 determined based upon the average conversion factor for freshwater (FW) and saltwater wetlands (SW).
12 The used annual flux was the measured annual flux if that was available; otherwise, it was the calculated
13 annual flux.

14
15 In cases where multiple years from the same site were presented, we took the average of those years.
16 Similarly, when multiple sites of the same type were presented in the same paper, we took the average.
17 Studies were separated into freshwater and estuarine systems.

18 In cases where papers presented both an annual flux and a mean daily flux, we calculated a
19 conversion factor [annual flux/(average daily flux $\times 10^3$)] to quantify the relationship between those two
20 numbers (Table 13A-5). When we looked at all studies ($n = 30$), this conversion factor was 0.36,
21 suggesting that there is a 360-day emission season. There was surprisingly little variation in this ratio, and
22 it was similar in freshwater (0.36) and estuarine (0.34) wetlands. In contrast, previous syntheses used a
23 150-day emission season for temperate wetlands (Matthews and Fung, 1987, Bartlett and Harriss, 1993).
24 While substantial winter methane emissions have been found in some studies, it is likely that flux data
25 from most studies have a non-normal distribution with occasional periods of high flux rates that are better
26 captured with annual measurements.

27 Using the conversion factors for freshwater and estuarine wetlands, we estimated average annual
28 fluxes from the average daily fluxes. For freshwater wetlands, the calculated average annual flux rate was
29 $38.6 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$ ($n = 74$), which is slightly larger than the average actual measured flux rate of
30 $32.1 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$ ($n = 32$). For estuarine wetlands, the average calculated annual flux rate was
31 $9.8 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$ ($n = 25$), which is smaller than the average measured flux rate of $16.9 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$
32 ($n = 13$). However, if we remove one outlier, the average measured flux rate is $10.2 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$.

33 Finally, we combined both approaches. In cases where a paper presented an annual value, we used
34 that number. In cases where only an average daily number was presented, we used that value corrected

1 with the appropriate conversion factor. For conterminous U.S. wetlands, FWMS Canadian wetlands, and
2 Mexican wetlands, we used an average flux of $36 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$, and for estuarine wetlands, we used an
3 average flux of $10.3 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$.

5 Plant Carbon Fluxes

6 We have limited our focus on plant carbon fluxes to those processes that would result in the
7 accumulation of plant carbon biomass on an interannual basis. Tree biomass carbon sequestration
8 averages $-140 \text{ g C m}^{-2} \text{ yr}^{-1}$ in U.S. forests across all forest types (Birdsey, 1992). Using the tree growth
9 estimates from the southeastern U.S. regional assessment of wetland forests (Brown *et al.*, 2001) yields an
10 even lower estimate of sequestration in aboveground tree biomass (approx. $-50.2 \text{ g C m}^{-2} \text{ yr}^{-1}$). We have
11 used this lower value to estimate that U.S. wetland forests currently sequester $-10.3 \text{ Mt C yr}^{-1}$.

12 We have assumed that the largely undisturbed forested wetlands of Alaska and Canada are at an
13 approximate steady state in terms of biomass, with no interannual plant carbon accumulation. It is likely
14 that plant carbon sequestration occurs largely as woody biomass, so we also assumed that non-forested
15 wetlands have no interannual plant carbon accumulation.

18 REFERENCES

- 19 Alford, D. P., R. D. Delaune, and C. W. Lindau, 1997: Methane flux from Mississippi River deltaic plain wetlands.
20 *Biogeochemistry*, **37**, 227-236.
- 21 Armentano, T. B., and E. S. Menges, 1986: Patterns of change in the carbon balance of organic soil- wetlands of the
22 temperate zone. *Journal of Ecology*, **74**, 755-774.
- 23 Aselmann, I., and P. J. Crutzen, 1989: Global distribution of natural freshwater wetlands and rice paddies, their net
24 primary productivity, seasonality and possible methane emissions. *Journal of Atmospheric Chemistry*, **8**, 307-
25 359.
- 26 Bartlett, K. B., and R. C. Harriss, 1993: Review and assessment of methane emissions from wetlands. *Chemosphere*,
27 **26**, 261-320.
- 28 Batjes, N. H., 1996: Total carbon and nitrogen in the soils of the world. *European Journal of Soil Science*, **47**, 151-
29 163.
- 30 Birdsey, R. A., 1992: Carbon storage and accumulation in United States forest ecosystems. General Technical Report
31 WO-59 Washington, DC: USDA Forest Service.
- 32 Bridgham, S. D., C.-L. Ping, J. L. Richardson, and K. Updegraff, 2000: Soils of northern peatlands: Histosols and
33 Gelisols. In: *Wetland Soils: Genesis, Hydrology, Landscapes, and Classification* (J. L. Richardson, and M. J.
34 Vepraskas, eds.), 343-370. Boca Raton, FL: CRC Press.
- 35 Brown, M. J., G. M. Smith, and J. McCollum, 2001: Wetland forest statistics for the south Atlantic states. RB-SRS-
36 062 Asheville, North Carolina: Southern Research Station, U.S. Forest Service.

- 1 Chmura, G. L., S. C. Anisfeld, D. R. Cahoon, and J. C. Lynch, 2003: Global carbon sequestration in tidal, saline
2 wetland soils. *Global Biogeochemical Cycles*, **17**, 1111.
- 3 Cleary, J., N. T. Roulet, and T. R. Moore, 2005: Greenhouse gas emissions from Canadian peat extraction, 1990-
4 2000: A life-cycle analysis. *Ambio*, **34**, 456-461.
- 5 Clymo, R. S., J. Turunen, and K. Tolonen, 1998: Carbon accumulation in peatland. *Oikos*, **81**, 368-388.
- 6 Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe, 1979: Classification of wetlands and deepwater habitats
7 of the United States. FWS/OBS-79/31 Washington, DC: Fish and Wildlife Service, US Department of the
8 Interior.
- 9 Dahl, T. E., 1990: Wetland losses in the United States 1970's to 1980's. Washington, DC: Fish and Wildlife Service,
10 U.S. Department of the Interior.
- 11 Dahl, T. E., 2000: Status and Trends of Wetlands in the Conterminous United States, 1986 to 1997. Washington, DC:
12 Fish and Wildlife Service, U.S. Department of the Interior.
- 13 Dahl, T. E., and C. E. Johnson, 1991: Status and Trends of Wetlands in the Conterminous United States, Mid-1970's
14 to Mid-1980's. Washington, D.C.: U.S. Department of the Interior, Fish and Wildlife Service.
- 15 Davidson, I., R. Vanderkam, and M. Padilla, 1999: Review of wetland inventory information in North
16 America. Canberra, Australia.
- 17 Ehhalt, D., M. Prather, F. Dentener, E. Dlugokencky, E. Holland, I. Isaksen, J. Katima, V. Kirchhoff, P. Matson, P.
18 Midgley, and M. Wang, 2001: "Atmospheric chemistry and greenhouse gases." In *Climate Change 2001: The
19 Scientific Basis*. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental
20 Panel on Climate Change (J. T. Houghton, Y. Ding, D. J. Griggs, M. Noguer, P. J. van der Linden, X. Dai, K.
21 Maskell, and C. A. Johnson, eds.), 239-287. Cambridge: Cambridge University Press.
- 22 Eswaran, H., E. Van Den Berg, and J. Kimble, 1995: Global soil carbon resources. In: *Soils and Global Change* (R.
23 Lal, J. Kimble, E. Levine, and B. A. Stewart, eds.), 27-43. Boca Raton, Florida: Lewis Publishers.
- 24 FAO, 1991: The Digitized Soil Map of the World. World Soil Resource Report, 64 Rome: Food and Agriculture
25 Organization.
- 26 FAO-UNESCO, 1974: *Soil Map of the World (1:5,000,000)*. Paris: UNESCO.
- 27 Frayer, W. E., T. J. Monahan, D. C. Bowden, and F. A. Graybill, 1983: Status and Trends of Wetlands and
28 Deepwater Habitats in the Conterminous United States, 1950s to 1970s. Fort Collins, Colorado: Dept. of Forest
29 and Wood Sciences, Colorado State University.
- 30 Gorham, E., 1991: Northern peatlands: Role in the carbon cycle and probable responses to climatic warming.
31 *Ecological Applications*, **1**, 182-195.
- 32 Group, N. W. W., 1988: Wetlands of Canada: Sustainable Development Branch, Environment Canada, Ontario, and
33 Polyscience Publications, Montreal, Quebec.
- 34 Hall, J. V., W. E. Frayer, and B. O. Wilen, 1994: Status of Alaska Wetlands. Anchorage, Alaska: U.S. Fish and
35 Wildlife Service.
- 36 Halsey, L. A., D. H. Vitt, and L. D. Gignac, 2000: *Sphagnum*-dominated peatlands in North America since the last
37 glacial maximum: their occurrence and extent. *The Bryologist*, **103**, 334-352.

- 1 Hanson, A. R., and L. Calkins, 1996: Wetlands of the Maritime Provinces: Revised Documentation for the Wetlands
2 Inventory. Technical Report No. 267 Sackville, New Brunswick: Canadian Wildlife Service, Atlantic Region.
- 3 Johnston, C. A., 1991: Sediment and nutrient retention by freshwater wetlands: effects on surface water quality.
4 *Critical Reviews in Environmental Control*, **21**, 491-565.
- 5 Joosten, H., and D. Clarke, 2002: Wise Use of Mires and Peatlands - Background Principles including a Framework
6 for Decision-Making. Saarijärvi, Finland: International Mire Conservation Group and International Peat
7 Society.
- 8 Kelly, C. A., J. W. M. Rudd, R. A. Bodaly, N. T. Roulet, V. L. St. Louis, A. Heyes, T. R. Moore, S. Schiff, R.
9 Aravena, K. J. Scott, B. Dyck, R. Harris, B. Warner, and G. Edwards, 1997: Increase in fluxes of greenhouse
10 gases and methyl mercury following flooding of an experimental reservoir. *Environmental Science &*
11 *Technology*, **31**, 1334-1344.
- 12 Kivinen, E., and P. Pakarinen, 1981: Geographical distribution of peat resources and major peatland complex types
13 in the world. *Annales Academiae Scientiarum Fennicae, Series A. III.* **132**, 1-28.
- 14 Lappalainen, E., 1996: General review on world peatland and peat resources. In: *Global Peat Resources* (E.
15 Lappalainen, ed.), 53-56. Jyskä, Finland: International Peat Society and Geological Survey of Finland.
- 16 Maltby, E., and P. Immirzi, 1993: Carbon dynamics in peatlands and other wetland soils, regional and global
17 perspectives. *Chemosphere*, **27**, 999-1023.
- 18 Malterer, T. J., 1996: Peat resources of the United States. In: *Global Peat Resources* (E. Lappalainen, ed.), 253-260.
19 Jyska, Finland.
- 20 Matthews, E., and I. Fung, 1987: Methane emission from natural wetlands: Global distribution, area, and
21 environmental characteristics of sources. *Global Biogeochemical Cycles*, **1**, 61-86.
- 22 Meade, R. H., T. R. Yuzyk, and T. J. Day, 1990: Movement and storage of sediments in rivers of the United States
23 and Canada. In: *Surface Water Hydrology, Geol. of N. Am., 0-1* (M. G. Wolman, and H. C. Riggs, eds.), 255-
24 280. Boulder, CO: Geological Society of American.
- 25 Mendelsohn, I. A., and K. L. McKee, 2000: Saltmarshes and mangroves. In: *North American Terrestrial Vegetation*
26 (M. G. Barbour, and W. D. Billings, eds.), 501-536. Cambridge, UK: Cambridge University Press.
- 27 Mitsch, W. J., and J. G. Gosselink, 1993: Wetlands New York: Van Nostrand Reinhold.
- 28 Moore, T. R., and N. T. Roulet, 1995: Methane emissions from Canadian peatlands. In: *Soils and Global Change* (R.
29 Lal, J. Kimble, E. Levine, and B. A. Stewart, eds.), 153-164. Boca Raton, FL: Lewis Publishers.
- 30 Moore, T. R., N. T. Roulet, and J. M. Waddington, 1998: Uncertainty in predicting the effect of climatic change on
31 the carbon cycling of Canadian peatlands. *Climatic Change*, **40**, 229-245.
- 32 National Wetlands Working Group, 1988: *Wetlands of Canada*, Sustainable Development Branch, Environment
33 Canada, Ontario, and Polyscience Publications, Montreal, Quebec.
- 34 NRCS, 1999: Soil Taxonomy: A Basic System of Soil Classification for Making and Interpreting Soil Surveys.
35 Washington, DC: Natural Resources Conservation Service, U.S. Department of Agriculture.
- 36 Ovenden, L., 1990: Peat accumulation in northern wetlands. *Quaternary Research*, **33**, 377-386.

- 1 Rieger, S., D. B. Schoepfoster, and C. E. Furbush, 1979: Exploratory Soil Survey of Alaska. Anchorage, Alaska:
2 USDA Soil Conservation Service.
- 3 Robinson, S. D., and T. R. Moore, 1999: Carbon and peat accumulation over the past 1200 years in a landscape with
4 discontinuous permafrost, northwestern Canada. *Global Biogeochemical Cycles*, **13**, 591-602.
- 5 Rubec, C., 1996: The status of peatland resources in Canada. In: *Global Peat Resources* (E. Lappalainen, ed., 243-
6 252. Jyskä, Finland: International Peat Society and Geological Survey of Finland.
- 7 Spalding, M., F. Blasco, and C. Field, eds., 1997: World Mangrove Atlas. Okinawa, Japan: The International
8 Society for Mangrove Ecosystems.
- 9 Spiers, A. G., 1999: Review of international/continental wetland resources. In: *Global Review of Wetland Resources*
10 *and Priorities* (C. M. Finlayson, and A. G. Spiers, eds.).
- 11 Stallard, R. F., 1998: Terrestrial sedimentation and the carbon cycle: Coupling weathering and erosion to carbon
12 burial. *Global Biogeochemical Cycles*, **12**, 231–257.
- 13 Tarnocai, C., 1998: The amount of organic carbon in various soil orders and ecological provinces in Canada. In: *Soil*
14 *Processes and the Carbon Cycle* (R. Lal, J. M. Kimble, R. F. Follett, and B. A. Stewart, eds.), 81-92. Boca
15 Raton, Florida: CRC Press.
- 16 Tarnocai, C., I. M. Kettles, and B. Lacelle, 2005: Peatlands of Canada. Ottawa.Ottawa, Canada: Agriculture and
17 Agri-Food Canada, Research Branch.
- 18 Tolonen, K., and J. Turunen, 1996: Accumulation rates of carbon in mires in Finland and implications for climactic
19 change. *Holocene*, **6**, 171-178.
- 20 Trumbore, S. E., and J. W. Harden, 1997: Accumulation and turnover of carbon in organic and mineral soils of the
21 BOREAS northern study area. *Journal of Geophysical Research*, **102**, 28, 817-828, 830.
- 22 Turetsky, M. R., R. K. Wieder, L. A. Halsey, and D. Vitt, 2002: Current distribution and diminishing peatland
23 carbon sink. *Geophysical Research Letters*, **29**, 10.1029/2001GL014000, 012002.
- 24 Turunen, J., N. T. Roulet, and T. R. Moore, 2004: Nitrogen deposition and increased carbon accumulation in
25 ombrotrophic peatlands in eastern Canada. *Global Biogeochemical Cycles*, **18**, GB3002,
26 doi:3010.1029/2003GB002154.
- 27 Twilley, R. R., R. H. Chen, and T. Hargis, 1992: Carbon sinks in mangroves and their implications to carbon budget
28 of tropical coastal ecosystems. *Water, Air and Soil Pollution*, **64**, 265-288.
- 29 Vitt, D. H., L. A. Halsey, I. E. Bauer, and C. Campbell, 2000: Spatial and temporal trends in carbon storage of
30 peatlands of continental western Canada through the Holocene. *Canadian Journal of Earth Sciences*, **37**, 683-
31 693.
- 32 Vitt, D. H., L. A. Halsey, and S. C. Zoltai, 1994: The bog landforms of continental western Canada in relation to
33 climate and permafrost patterns. *Arctic and Alpine Research*, **26**, 1-13.
- 34 Webb, R. S., and T. Webb III, 1988: Rates of sediment accumulation in pollen cores from small lakes and mires of
35 eastern North America. *Quaternary Research*, **30**, 284-297.
- 36 WEC, 2001: Survey of Energy Resources. [http://www.worldenergy.org/wec-](http://www.worldenergy.org/wec-geis/publications/reports/ser/peat/peat.asp)
37 [geis/publications/reports/ser/peat/peat.asp](http://www.worldenergy.org/wec-geis/publications/reports/ser/peat/peat.asp)

- 1 Werner, C., K. Davis, P. Bakwin, C. Yi, D. Hurst, and L. Lock, 2003: Regional-scale measurements of CH₄
2 exchange from a tall tower over a mixed temperate/boreal lowland and wetland forest. *Global Change Biology*,
3 **9**, 1251-1261.
- 4 West, A. E., P. D. Brooks, M. C. Fisk, L. K. Smith, E. A. Holland, C. H. Jaeger III, S. Babcock, R. S. Lai, and S. K.
5 Schmidt, 1999: Landscape patterns of CH₄ fluxes in an alpine tundra ecosystem. *Biogeochemistry*, **45**, 243-264.
- 6 Wickland, K. P., R. G. Striegl, S. K. Schmidt, and M. A. Mast, 1999: Methane flux in subalpine wetland and
7 unsaturated soils in the southern Rocky Mountains. *Global Biogeochemical Cycles*, **13**, 101-113.
- 8 Wilson, J. O., P. M. Crill, K. B. Bartlett, D. I. Sebacher, R. C. Harriss, and R. L. Sass, 1989: Seasonal variation of
9 methane emissions from a temperate swamp. *Biogeochemistry*, **8**, 55-71.
- 10 Yavitt, J. B., 1997: Methane and carbon dioxide dynamics in *Typha latifolia* (L.) wetlands in central New York
11 state. *Wetlands*, **17**, 394-406.
- 12 Yavitt, J. B., G. E. Lang, and A. J. Sexstone, 1990: Methane fluxes in wetland and forest soils, beaver ponds, and
13 low-order streams of a temperate forest ecosystem. *Journal of Geophysical Research*, **95**, 22463-22474.
- 14 Yavitt, J. B., R. K. Wieder, and G. E. Lang, 1993: CO₂ and CH₄ dynamics of a *Sphagnum*-dominated peatland in
15 West Virginia. *Global Biogeochemical Cycles*, **7**, 259-274.

1 **Table 13A-1. Current and historical area of wetlands in North America and the world ($\times 10^3$ km²).** Historical refers to approximately 1800, unless otherwise
 2 specified.

	Permafrost peatlands	Non-permafrost peatlands	Mineral-soil freshwater	Salt marsh	Mangrove	Mudflat	Total
<u>Canada</u>							
Current	422 ^a	714 ^a	159 ^b	0.4 ^c	0	6 ^d	1301
Historical	424 ^e	726 ^f	359 ^g	1.3 ^b	0	7 ^h	1517
<u>Alaska</u>							
Current	89 ⁱ	43 ⁱ	556 ^j	1.4 ^c	0	7 ^k	696
Historical	89	43	556	1.4	0	9 ^b	698
<u>Conterminous United States</u>							
Current	0	93 ^L	312 ^m	18 ^c	3 ^c	2 ⁿ	428
Historical	0	111 ⁱ	762 ^o	20 ^p	4 ⁿ	3 ⁿ	899
<u>Mexico</u>							
Current	0	10 ^p	21 ^q	0	5 ^c	ND ^r	36
Historical	0		45 ^p	0	7 ^h	ND	52
<u>North America</u>							
Current	511	861	1,047	20	8	15	2,461
Historical	513	894 ^s	1,706 ^s	23	11	19	3,166
<u>Global</u>							
Current	3,443 ^t		2,289 to 2,341 ^u	22 ^v	181 ^w	ND	~6,000
Historical	3,880-4,086 ^x		5,000 ^y	ND	ND	ND	~9,000 ^y

3
 4 ^aTarnocai *et al.* (2005).

5 ^bNational Wetlands Working Group (1988).

6 ^cMendelssohn and McKee (2000).

7 ^dEstimated from the area of Canadian salt marshes and the ratio of mudflat to salt marsh area reported by Hanson and Calkins (1996).

8 ^eAccounting for losses due to permafrost melting in western Canada (Vitt *et al.*, 1994). This is an underestimate, as similar, but undocumented, losses have
 9 probably also occurred in eastern Canada and Alaska.

10 ^f9000 km² lost to reservoir flooding (Rubec, 1996), 250 km² to forestry drainage (Rubec, 1996), 124 km² to peat harvesting for horticulture (Cleary *et al.*,
 11 2005), and 16 km² to oil sands mining (Turetsky *et al.*, 2002). See note e for permafrost melting estimate.

- 1 ^gRubec (1996).
- 2 ^hAssumed same loss rate as the conterminous United States since 1954 (Dahl, 2000).
- 3 ⁱHistorical area from NRCS soil inventory (Bridgham *et al.*, 2000), except Alaska inventory updated by N. Bliss from a February 2006 query of the
4 STATSGO database. Less than 1% wetland losses have occurred in Alaska (Dahl, 1990).
- 5 ^jTotal freshwater wetland area from Hall *et al.* (1994) minus peatland area.
- 6 ^kHall *et al.*, 1994.
- 7 ^LHistorical area from Bridgham *et al.* (2000) minus losses in Armentano and Menges (1986).
- 8 ^mOverall freshwater wetland area from Dahl (2000) minus peatland area.
- 9 ⁿDahl (2000).
- 10 ^oTotal historical wetland area from Dahl (1990) minus historical peatland area minus historical estuarine area.
- 11 ^pDavidson *et al.* (1999).
- 12 ^qSpiers (1999).
- 13 ^rND indicates that no data are available.
- 14 ^sAssuming that historical proportion of peatlands to total wetlands in Mexico was the same as today.
- 15 ^tBridgham *et al.* (2000) for the United States, Tarnocai *et al.* (2005) for Canada, Joosten and Clarke (2002) for the rest of world. Recent range in literature
16 2,974,000–3,985,000 km² (Matthews and Fung, 1987; Aselmann and Crutzen, 1989; Maltby and Immerzi, 1993; Bridgham *et al.*, 2000; Joosten and Clarke,
17 2002).
- 18 ^uMatthews and Fung (1987); Aselmann and Crutzen (1989). For subsequent calculations, used the average of 2,315,000 km².
- 19 ^vChmura *et al.* (2003). Underestimated because no inventories were available for the continents Asia, South America and Australia which are mangrove-
20 dominated but also support salt marsh.
- 21 ^wSpalding *et al.* (1997).
- 22 ^xMaltby and Immerzi (1993). For subsequent calculations, used 4,000,000 km².
- 23 ^yApproximately 50% loss from Moser *et al.* (1996). For subsequent calculations, used an original global mineral-soil wetland area of 5,000,000 km².

1 **Table 13A-2. Soil carbon pools (Gt) and fluxes (Mt yr⁻¹) of wetlands in North America and the world.** “Sequestration in current wetlands” refers to carbon
 2 sequestration in wetlands that currently exist; “oxidation in former wetlands” refers to emissions from wetlands that have been converted to non-wetland uses or
 3 conversion among wetland types due to human influence; “historical loss in sequestration capacity” refers to the loss in the carbon sequestration function of
 4 wetlands that have been converted to non-wetland uses; “change in flux from wetland conversions” is the sum of the two previous fluxes. Positive flux numbers
 5 indicate a net flux into the atmosphere, whereas negative numbers indicate a net flux into the ecosystem
 6

	Permafrost peatlands	Non-permafrost peatlands	Mineral-soil freshwater	Salt marsh	Mangrove	Mudflat	Total
<u>Canada</u>							
Pool Size in Current Wetlands	44.2 ^a	102.9 ^a	4.6 ^b	0.0 ^c	0.0	0.1 ^d	151.8
Sequestration in Current Wetlands	-5.5 ^e	-13.6 ^f	-5.1 ^g	-0.1	0.0	-1.2 ^d	-25.5
Oxidation in Former Wetlands		0.2 ^h	0.0 ⁱ	0.0 ^j	0.0	0.0	0.2
Historical Loss in Sequestration Capacity	0.0 ^e	0.2 ^f	6.5 ^g	0.2	0.0	0.3	7.2
Change in Flux From Wetland Conversions		0.4	6.5	0.2	0.0	0.3	7.4
<u>Alaska</u>							
Pool Size in Current Wetlands	9.3 ^k	6.2 ^k	26.0 ^L	0.0	0.0	0.1	41.7
Sequestration in Current Wetlands	-1.1 ^e	-0.8 ^f	-18.0 ^g	-0.3	0.0	-1.6	-21.9
Oxidation in Former Wetlands	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Historical Loss in Sequestration Capacity	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Change in Flux From Wetland Conversions	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<u>Conterminous United States</u>							
Pool Size in Current Wetlands	0	14.0 ^m	5.1 ^L	0.4	0.1	0.1	19.7
Sequestration in Current Wetlands	0	-11.6 ⁿ	-10.1 ^g	-3.9	-0.5	-0.5	-26.6
Oxidation in Former Wetlands	0	18.0 ^o	0.1 ⁱ	0.0	0.0	0.0	18.1
Historical Loss in Sequestration Capacity	0	2.1 ⁿ	14.5 ^g	0.3	0.0	0.1	17.1
Change in Flux from Wetland Conversions	0	20.1	14.6	0.3	0.0	0.1	35.2
<u>Mexico</u>							
Pool Size in Current Wetlands	0.0	1.5 ^m	0.3 ^L	0.0	0.1	ND*	1.9

Sequestration in Current Wetlands	0	-1.6 ^p	-0.7 ^g	0.0	-1.6	ND	-3.9
Oxidation in Former Wetlands	0	ND	ND	0.0	0.0	0.0	ND
Historical Loss in Sequestration Capacity	0	ND	ND	0.0	0.5	ND	0.5
Change in Flux from Wetland Conversions	0	ND	ND	0.0	0.5	ND	0.5
North America							
Pool Size in Current Wetlands	53.5	124.6	36.0	0.4	0.2	0.3	215.1
Sequestration in Current Wetlands	-6.6	-27.6	-33.9	-4.3	-2.1	-3.3	-77.8
Oxidation in Former Wetlands	18.2		0.1	0.0	0.0	0.0	18.3
Historical Loss in Sequestration Capacity	0	2.3	21.0	0.5	0.5	0.5	24.8
Change in Flux from Wetland Conversions	20.5		21.1	0.5	0.5	0.5	43.1
Global							
Pool Size in Current Wetlands	234 to 679 ^q		46 ^r	0.4 ^s	5.0 ^s	ND	286 to 730
Sequestration in Current Wetlands	-40 to -70 ^t		-75 ^g	-4.6 ^s	-38.0 ^s	ND	-158 to -188
Oxidation in Former Wetlands	160 to 250 ^u		ND	0	0	0	160 to 250
Historical Loss in Sequestration Capacity	16 ^u		87 ^g	0.5 ^v	12.7 ^w	ND	116
Change in Flux From Wetland Conversions	176 to 266 ^u		> 87 ^x	0.5	12.7	ND	276 to 366

- 1
- 2 *ND indicates that no data are available.
- 3 ^aTarnocai *et al.* (2005).
- 4 ^bTarnocai (1998).
- 5 ^cRates calculated from Chimura *et al.* (2003); areas from Mendelssohn and McKee (2000).
- 6 ^dAssumed the same carbon density and accumulation rates as the adjacent vegetated wetland ecosystem (mangrove data for Mexico and salt marsh data
- 7 elsewhere).
- 8 ^eSoil carbon accumulation rate of 0.13 Mg C ha⁻¹ yr⁻¹ (see Chapter 12 in this report).
- 9 ^fCarbon accumulation rate of 0.19 Mg C ha⁻¹ yr⁻¹. This is an average value of the reported range of long-term apparent accumulation rate of 0.05–0.35
- 10 (Ovenden, 1990, Maltby and Immirzi, 1993; Trumbore and Harden, 1997; Vitt *et al.*, 2000; Turunen *et al.*, 2004).

- 1 ^gPotential rate calculated as the average sediment accumulation rate of 1680 g m⁻² yr⁻¹ (range 0–7840) from Johnston (1991) times 7.7% C (CV = 109) (Batjes,
2 1996). It was assumed that the actual rate was 25% of the potential rate because of bias in choosing sampling sites and considerations of the redistribution of
3 sediment due to erosion without a change in the sequestration rate on a landscape scale.
- 4 ^hSum of -0.24 Mt C yr⁻¹ from horticulture removal of peat (Cleary *et al.*, 2005) and 0.10 Mt C yr⁻¹ from increased peat sequestration due to permafrost melting
5 (Turetsky *et al.*, 2002).
- 6 ⁱAssumed that the oxidized soil C is lost over 50 yr.
- 7 ^jAssumed that conversion of tidal systems is caused by fill and results in burial and preservation of SOM define SOM rather than oxidation.
- 8 ^kSoil carbon densities of 1,441 Mg C ha⁻¹ for Histosols and 1,048 Mg C ha⁻¹ for Histels (Tarnocai *et al.*, 2005).
- 9 ^lSoil carbon density of 162 Mg C ha⁻¹ for the conterminous United States and Mexico and 468 Mg C ha⁻¹ for Alaska based upon NRCS STATSGO database
10 and soil pedon information.
- 11 ^mAssumed soil carbon density of 1,500 Mg C ha⁻¹.
- 12 ⁿWebb and Webb (1988).
- 13 ^oEstimated loss rate as of early 1980s (Armentano and Menges,1986). Overall wetlands losses in the United States have declined dramatically since then
14 (Dahl, 2000) and probably even more so for Histosols, so this number may still be representative.
- 15 ^pUsing peat accumulation rate of 1.6 Mg C ha⁻¹ (range 1.0–2.25) (Maltby and Immirzi, 1993).
- 16 ^qGorham (1991), Maltby and Immirzi (1993), Eswaran *et al.* (1995), Batjes (1996), Lappalainen (1996), Joosten and Clarke (2002).
- 17 ^rSoil carbon density of 199 Mg C ha⁻¹ (Batjes, 1996).
- 18 ^sChmura *et al.* (2003).
- 19 ^tJoosten and Clarke (2002). Using the peatland estimate in Table 13A-1 and a carbon accumulation rate of 0.19 Mg C ha⁻¹ yr⁻¹, we calculate a global flux of
20 –65 Mt C yr⁻¹ in peatlands.
- 21 ^uCurrent oxidative flux is the difference between the change in flux and the historical loss in sequestration capacity from this table. The change in flux is from
22 Maltby and Immirzi (1993) and the historical loss in sequestration capacity is from this table for North America, from Armentano and Menges (1986) for other
23 northern peatlands, and from Maltby and Immirzi (1993) for tropical peatlands.
- 24 ^vAssumed that global rates approximate the North America rate because most salt marshes inventoried are in North America.
- 25 ^wAssumed 25% loss globally since the late 1800s.
- 26 ^x> sign indicates that this a minimal loss estimate.

1 **Table 13A-3. Plant carbon pools (Gt) and fluxes (Mt yr⁻¹) of wetlands in North America and the world.** Positive flux numbers indicate a net
 2 flux into the atmosphere, whereas negative numbers indicate a net flux into the ecosystem.

	Permafrost peatlands	Non-perma- frost peatlands	Mineral- soil freshwater	Salt marsh	Mangrove	Total
<u>Canada</u>						
Pool Size in Current Wetlands		1.4 ^a	0.3 ^b	0.0 ^c	0.0	1.7
Sequestration in Current Wetlands	0.0	ND*		0.0	0.0	0.0
<u>Alaska</u>						
Pool Size in Current Wetlands		0.4 ^a	1.1 ^d	0.0	0.0	1.5
Sequestration in Current Wetlands	0.0	0.0	0.0	0.0	0.0	0.0
<u>Conterminous United States</u>						
Pool Size in Current Wetlands	0.0	1.5 ^d		0.0	0.0	1.5
Sequestration in Current Wetlands	0.0	-10.3 ^e		0.0	0.0	-10.3
<u>Mexico</u>						
Pool Size in Current Wetlands	0.0	0.0 ^b	0.0 ^b	0.0	0.1	0.1
Sequestration in Current Wetlands	0.0	ND	ND	0.0	ND	0.0
<u>North America</u>						
Pool Size in Current Wetlands	4.8			0.0	0.1	4.9
Sequestration in Current Wetlands	0.0	-10.3		0.0	ND	-10.3
<u>Global</u>						
Pool Size in Current Wetlands	6.9 ^b		4.6 ^b	0.0 ^f	4.0 ^g	15.5
Sequestration in Current Wetlands	0.0	ND	ND	0.0	ND	ND

3 *ND indicates that no data are available.
 4 ^aBiomass for non-forested peatlands from Vitt *et al.* (2000), assuming 50% of biomass is belowground. Forest biomass density from
 5 Birdsey (1992) and forested area from Tarnocai *et al.* (2005) for Canada and from Hall *et al.* (1994) for Alaska.
 6 ^bAssumed 2000 g C m⁻² in aboveground and belowground plant biomass (Gorham, 1991).
 7 ^cBiomass data from Mitsch and Gosselink (1993).
 8 ^dBiomass for non-forested wetlands from Gorham (1991). Forest biomass density from Birdsey (1992), and forested area from Dahl (2000).
 9 ^e50 g C m⁻² yr⁻¹ sequestration from forest growth from a southeastern U.S. regional assessment of wetland forest growth (Brown *et al.*, 2001).
 10 ^fAssumed that global pools approximate those from North America because most salt marshes inventoried are in North America.
 11 ^gTwilley *et al.* (1992).

1 **Table 13A-4. Methane fluxes (Mt yr⁻¹) from wetlands in North America and the world.**

	Permafrost peatlands	Non-permafrost peatlands	Mineral-soil freshwater	Salt marsh	Mangrove	Mudflat	Total
<u>Canada</u>							
CH ₄ Flux in Current Wetlands	1.1 ^a	2.1 ^{a,b}	5.7	0.0	0.0	0.0 ^c	8.9
Historical change in CH ₄ Flux	0.0	0.3	-7.2	0.0	0.0	0.0	-6.9
<u>Alaska</u>							
CH ₄ Flux in Current Wetlands	0.2	0.1	1.4	0.0	0.0	0.1	1.8
Historical change in CH ₄ Flux	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<u>Conterminous United States</u>							
CH ₄ Flux in Current Wetlands	0.0	3.4	11.2	0.1	0.0	0.0	14.7
Historical change in CH ₄ Flux	0.0	-0.6	-16.2	0.0	0.0	0.0	-16.8
<u>Mexico</u>							
CH ₄ Flux in Current Wetlands	0.0	0.4	0.7	0.0	0.0	ND*	1.1
Historical change in CH ₄ Flux	0.0	-0.5		0.0	0.0	ND	-0.5
<u>North America</u>							
CH ₄ Flux in Current Wetlands	1.3	5.9	19.1	0.1	0.1	0.1	26.5
Historical change in CH ₄ Flux	0.0	-24.2		0.0	0.0	0.0	-24.2
<u>Global</u>							
CH ₄ Flux in Current Wetlands	14.1 ^d	22.5 ^d	68.0 ^d	0.1 ^e	1.4	ND	92–237 ^f
Historical change in CH ₄ Flux	-3.6		-79	0.0 ^g	-0.5	ND	-83

2 *ND indicates that no data are available.

3 ^aUsed CH₄ flux of 2.5 g m⁻² yr⁻¹ (range 0 to 130, likely mean 2–3) (Moore and Roulet, 1995) for Canadian peatlands and all Alaskan freshwater wetlands. Used CH₄ flux of
 4 36.0 g m⁻² yr⁻¹ for Canadian freshwater mineral-soil wetlands and all U.S. and Mexican freshwater wetlands and 10.3 g m⁻² yr⁻¹ for estuarine wetlands—from synthesis of
 5 published CH₄ fluxes for the United States (see Table 13A-5).

6 ^bIncludes a 17-fold increase in CH₄ flux (Kelly *et al.*, 1997) in the 9000 km² of reservoirs that have been formed on peatlands (Rubec, 1996) and an estimated CH₄ flux of 15 g
 7 m⁻² yr⁻¹ (Moore *et al.*, 1998) from 2,630 km² of melted permafrost peatlands (Vitt *et al.*, 1994).

8 ^cAssumed trace gas fluxes from unvegetated estuarine wetlands (i.e., mudflats) was the same as adjacent wetlands.

9 ^dBartlett and Harriss (1993).

10 ^eAssumed that global rates approximate the North America rate because most salt marshes area is in North America.

11 ^fEhhalt *et al.* (2001).

12 ^gAssumed a conservative 25% loss since the late 1800s.

1 **Table 13A-5. Methane fluxes measured in the conterminous United States.** The conversion factor is the ratio of the daily average flux to the measured annual
 2 flux $\times 10^3$. The calculated annual flux was determined based upon the average conversion factor for freshwater (FW) and saltwater wetlands (SW). The used
 3 annual flux was the measured annual flux if that was available; otherwise, it was the calculated annual flux
 4

Habitat	State	Method ^a	Salt/ Fresh	Daily Average Flux (mg CH ₄ m ⁻² d ⁻¹)	Measured Annual Flux (g CH ₄ m ⁻² yr ⁻¹)	Conversion Factor	Calculated Annual Flux (g CH ₄ m ⁻² yr ⁻¹)	Used Annual Flux (g CH ₄ m ⁻² yr ⁻¹)	Reference
Fens	CO	C	FW		40.7			40.7	Chimner and Cooper (2003)
Wet Alpine Meadow	CO	C	FW	0.1			0.0	0.0	Neff <i>et al.</i> (1994)
Lake - Average	CO	C	FW	25.4			9.2	9.2	Smith and Lewis (1992)
Wetland - Average	CO	C	FW	28.3			10.3	10.3	Smith and Lewis (1992)
Nuphar Bed	CO	C	FW	202.1			73.6	73.6	Smith and Lewis (1992)
Tundra - Carex Meadow	CO	C	FW	2.8			1.0	1.0	West <i>et al.</i> (1999)
Tundra - Acomastylis Meadow	CO	C	FW	-0.5			-0.2	-0.2	West <i>et al.</i> (1999)
Tundra - Kobresia Meadow	CO	C	FW	-0.8			-0.3	-0.3	West <i>et al.</i> (1999)
Moist Grassy	CO	C	FW	6.1	1.9	0.32	2.2	1.9	Wickland <i>et al.</i> (1999)
Moist Mossy	CO	C	FW	1.5	0.5	0.33	0.5	0.5	Wickland <i>et al.</i> (1999)
Wetland	CO	C	FW		41.7			41.7	Wickland <i>et al.</i> (1999)
Hardwood Hammock	FL	C	FW	0.0			0.0	0.0	Bartlett <i>et al.</i> (1989)
Dwarf Cypress / Sawgrass	FL	C	FW	7.5			2.7	2.7	Bartlett <i>et al.</i> (1989)
Spikerush	FL	C	FW	29.4			10.7	10.7	Bartlett <i>et al.</i> (1989)
Sawgrass < 1m	FL	C	FW	38.8			14.1	14.1	Bartlett <i>et al.</i> (1989)
Sawgrass/Spkerush/Periphyton	FL	C	FW	45.1			16.4	16.4	Bartlett <i>et al.</i> (1989)
Swamp Forest	FL	C	FW	68.9			25.1	25.1	Bartlett <i>et al.</i> (1989)
Sawgrass > 1m	FL	C	FW	71.9			26.2	26.2	Bartlett <i>et al.</i> (1989)
Sawgrass	FL	C	FW	107.0			38.9	38.9	Burke <i>et al.</i> (1988)
Pond Open Water	FL	C	FW	624.0			227.1	227.1	Burke <i>et al.</i> (1988)
Everglades - Cladium	FL	C	FW	45.4			16.5	16.5	Chanton <i>et al.</i> (1993)
Everglades - Typha	FL	C	FW	142.9			52.0	52.0	Chanton <i>et al.</i> (1993)
Wet Prairie (Marl)	FL	C	FW	87.0			31.6	31.6	Happell <i>et al.</i> (1993)
Wet Prairie (Marl)	FL	C	FW	27.4			10.0	10.0	Happell <i>et al.</i> (1993)
Marsh (Marl)	FL	C	FW	30.0			10.9	10.9	Happell <i>et al.</i> (1993)
Marsh (Marl)	FL	C	FW	49.6			18.0	18.0	Happell <i>et al.</i> (1993)

Marsh (Peat)	FL	C	FW	45.4			16.5	16.5	Happell <i>et al.</i> (1993)
Marsh (Peat)	FL	C	FW	13.0			4.7	4.7	Happell <i>et al.</i> (1993)
Marsh (Peat)	FL	C	FW	163.6			59.6	59.6	Happell <i>et al.</i> (1993)
Marsh (Peat)	FL	C	FW	20.4			7.4	7.4	Happell <i>et al.</i> (1993)
Wet Prairie / Sawgrass	FL	C	FW	61.0			22.2	22.2	Harriss <i>et al.</i> (1988)
Wetland Forest	FL	C	FW	59.0			21.5	21.5	Harriss <i>et al.</i> (1988)
Cypress Swamp - Flowing Water	FL	C	FW	67.0			24.4	24.4	Harriss and Sebacher (1981)
Open Water Swamp	FL	C	FW	480.0			174.7	174.7	Schipper and Reddy (1994)
Waterlily Slough	FL	C	FW	91.0			33.1	33.1	Schipper and Reddy (1994)
Cypress Swamp - Deep Water	GA	C	FW	92.3			33.6	33.6	Harriss and Sebacher (1981)
Bottotmand Hardwoods/ Swamps	GA	C	FW		23.0			23.0	Pulliam (1993)
Swamp Forest	LA	C	FW	146.0			53.1	53.1	Alford <i>et al.</i> (1997)
Freshwater Marsh	LA	C	FW	251.0			91.4	91.4	Alford <i>et al.</i> (1997)
Fresh	LA	C	FW	587.0	213.0	0.36	213.6	213.0	DeLaune <i>et al.</i> (1983)
Fresh	LA	C	FW	49.0	18.7	0.38	17.8	18.7	DeLaune <i>et al.</i> (1983)
Sphagnum Bog	MD	C	FW	-1.1			-0.4	-0.4	Yavitt <i>et al.</i> (1990)
Bog	MI	C	FW	193.0			70.2	70.2	Shannon and White (1994)
Bog	MI	C	FW	28.0			10.2	10.2	Shannon and White (1994)
Beaver Meadow	MN	C	FW		2.3			2.3	Bridgham <i>et al.</i> (1995)
Open Bogs	MN	C	FW		0.0			0.0	Bridgham <i>et al.</i> (1995)
Bog (Forested Hummock)	MN	C	FW	10.0	3.5	0.35	3.6	3.5	Dise (1993)
Bog (Forested Hollow)	MN	C	FW	38.0	13.8	0.36	13.8	13.8	Dise (1993)
Fen Lagg	MN	C	FW	35.0	12.6	0.36	12.7	12.6	Dise (1993)
Bog (Open Bog)	MN	C	FW	118.0	43.1	0.37	42.9	43.1	Dise (1993)
Fen (Open Poor Fen)	MN	C	FW	180.0	65.7	0.37	65.5	65.7	Dise (1993)
Poor Fen	MN	C	FW	242.0			88.1	88.1	Dise and Verry (2001)
Sedge Meadow	MN	C	FW		11.7			11.7	Naiman <i>et al.</i> (1991)
Submergent	MN	C	FW		14.4			14.4	Naiman <i>et al.</i> (1991)
Deep Water	MN	C	FW		0.5			0.5	Naiman <i>et al.</i> (1991)
Poor Fen	MN	T	FW		14.6			14.6	Shurpali and Verma (1998)
Submerged Tidal	NC	C, E	FW	144.8			52.7	52.7	Kelley <i>et al.</i> (1995)
Banks Tidal	NC	C, E	FW	20.1			7.3	7.3	Kelley <i>et al.</i> (1995)
Tidal Marsh	NC	C	FW	3.0	1.0	0.34	1.1	1.0	Megonigal and Schlesinger (2002)
Tidal Marsh	NC	C	FW	3.5	2.3	0.65	1.3	2.3	Megonigal and Schlesinger (2002)
Prairie Marsh	NE	T	FW		64.0			64.0	Kim <i>et al.</i> (1998)
Poor Fen	NH	C	FW	503.3	110.6	0.22	183.2	110.6	Carroll and Crill (1997)

Poor Fen	NH	C	FW		69.3			69.3	Frolking and Crill (1994)
Forested Peatland	NY	C	FW	0.6	0.2	0.37	0.2	0.2	Coles and Yavitt (2004)
Pools Forested Swamp	NY	C	FW	224.6	69.0	0.31	81.7	69.0	Miller <i>et al.</i> (1999)
Typha Marsh - Mineral Soils	NY	C	FW	344.4			125.3	125.3	Yavitt (1997)
Typha Marsh - Peat Soils	NY	C	FW	65.1			23.7	23.7	Yavitt (1997)
Typha Marsh - All soils	NY	C	FW	204.8			74.5	74.5	Yavitt (1997)
Cypress Swamp - Floodplain	SC	C	FW	9.9			3.6	3.6	Harriss and Sebacher (1981)
Swamp	VA	C	FW	470.3			171.2	171.2	Chanton <i>et al.</i> (1992)
Maple/gum Forested Swamp	VA	C	FW		0.5			0.5	Harriss <i>et al.</i> (1982)
Emergent Tidal Freshwater Marsh	VA	C	FW		96.2			96.2	Neubauer <i>et al.</i> (2000)
Oak Swamp (Bank Site)	VA	C	FW	117.0	43.7	0.37	42.6	43.7	Wilson <i>et al.</i> (1989)
Emergent Macrophytes (Peltandra)	VA	C	FW	155.0			56.4	56.4	Wilson <i>et al.</i> (1989)
Emergent Macrophytes (Smartweed)	VA	C	FW	83.0			30.2	30.2	Wilson <i>et al.</i> (1989)
Ash Tree Swamp	VA	C	FW	152.0			55.3	55.3	Wilson <i>et al.</i> (1989)
Bog	WA	C	FW	73.0			26.6	26.6	Lansdown <i>et al.</i> (1992)
Lowland Shrub and Forested Wetland	WI	T	FW		12.4			12.4	Werner <i>et al.</i> (2003)
Sphagnum Eriophorum (Poor Fen)	WV	C	FW	6.6			2.4	2.4	Yavitt <i>et al.</i> (1990)
Sphagnum Shrub (Fen)	WV	C	FW	0.1			0.0	0.0	Yavitt <i>et al.</i> (1990)
Polytrichum Shrub (Fen)	WV	C	FW	-0.1			0.0	0.0	Yavitt <i>et al.</i> (1990)
Sphagnum Forest	WV	C	FW	9.6			3.5	3.5	Yavitt <i>et al.</i> (1990)
Sedge Meadow	WV	C	FW	1.5			0.5	0.5	Yavitt <i>et al.</i> (1990)
Beaver Pond	WV	C	FW	250.0			91.0	91.0	Yavitt <i>et al.</i> (1990)
Low Gradient Headwater Stream	WV	C	FW	300.0			109.2	109.2	Yavitt <i>et al.</i> (1990)
Sphagnum-Eriophorum	WV	C	FW	52.1	19.0	0.37	18.9	19.0	Yavitt <i>et al.</i> (1993)
Polytrichum	WV	C	FW	41.1	15.0	0.37	15.0	15.0	Yavitt <i>et al.</i> (1993)
Sphagnum-Shrub	WV	C	FW	4.4	1.6	0.37	1.6	1.6	Yavitt <i>et al.</i> (1993)
Salt Marsh	DE	C	SW	0.5			0.2	0.2	Bartlett <i>et al.</i> (1985)
Red Mangroves	FL	C	SW	4.2			1.4	1.4	Bartlett <i>et al.</i> (1989)
Dwarf Red Mangrove	FL	C	SW	81.9			27.9	27.9	Bartlett <i>et al.</i> (1989)
High Marsh	FL	C	SW	3.9			1.3	1.3	Bartlett <i>et al.</i> (1985)
Salt Marsh	FL	C	SW	0.6			0.2	0.2	Bartlett <i>et al.</i> (1985)
Salt Water Mangroves	FL	C	SW	4.0			1.4	1.4	Harriss <i>et al.</i> (1988)
Salt Marsh	GA	C	SW	13.4			4.6	4.6	Bartlett <i>et al.</i> (1985)
Short Spartina Marsh - High Marsh	GA	C	SW	145.2	53.1	0.37	49.5	53.1	King and Wiebe (1978)
Mid Marsh	GA	C	SW	15.8	5.8	0.37	5.4	5.8	King and Wiebe (1978)
Tall Spartina Marsh - Low Marsh	GA	C	SW	1.2	0.4	0.34	0.4	0.4	King and Wiebe (1978)

Intermediate Marsh	LA	C	SW	912 ^b					Alford <i>et al.</i> (1997)
Salt Marsh	LA	C	SW	15.7	5.7	0.36	5.4	5.7	DeLaune <i>et al.</i> (1983)
Brackish	LA	C	SW	267.0	97.0		91.1	97.0	DeLaune <i>et al.</i> (1983)
Salt Marsh	LA	C	SW	4.8	1.7	0.35	1.6	1.7	DeLaune <i>et al.</i> (1983)
Brackish	LA	C	SW	17.0	6.4	0.38	5.8	6.4	DeLaune <i>et al.</i> (1983)
Cypress Swamp - Floodplain	SC	C	SW	1.5			0.5	0.5	Bartlett <i>et al.</i> (1985)
Salt Marsh	SC	C	SW	0.4			0.1	0.1	Bartlett <i>et al.</i> (1985)
Salt Marsh	VA	C	SW	3.0	1.3	0.43	1.0	1.3	Bartlett <i>et al.</i> (1985)
Salt Marsh	VA	C	SW	5.0	1.2	0.24	1.7	1.2	Bartlett <i>et al.</i> (1985)
Salt Meadow	VA	C	SW	2.0	0.4	0.22	0.7	0.4	Bartlett <i>et al.</i> (1985)
Salt Marsh	VA	C	SW	-0.8			-0.3	-0.3	Bartlett <i>et al.</i> (1985)
Salt Marsh	VA	C	SW	1.5			0.5	0.5	Bartlett <i>et al.</i> (1985)
Salt Meadow	VA	C	SW	-1.9			-0.6	-0.6	Bartlett <i>et al.</i> (1985)
Tidal Salt Marsh	VA	C	SW	16.0	5.6	0.35	5.5	5.6	Bartlett <i>et al.</i> (1987)
Tidal Brackish Marsh	VA	C	SW	64.6	22.4	0.35	22.0	22.4	Bartlett <i>et al.</i> (1987)
Tidal Brackish/Fresh Marsh	VA	C	SW	53.5	18.2	0.34	18.2	18.2	Bartlett <i>et al.</i> (1987)

FW

Average = 32.1 0.36 38.6 36.0

FW n = 32 18 74 88

FW

StError= 7.9 0.02 6.0 5.0

SW

Average = 16.9 0.34 9.8 10.3

SW n = 13 12 25 25

SW

StError= 7.8 0.02 4.1 4.4

1

2 ^aC = chamber, T = tower, eddy covariance, E = ebullition measured separately.

3 ^bOutlier that was removed from further analysis.

1

[This page intentionally left blank]